

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07F 9/6561, A61K 31/675		A1	(11) International Publication Number: WO 98/39344
			(43) International Publication Date: 11 September 1998 (11.09.98)
(21) International Application Number: PCT/US98/04502		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 6 March 1998 (06.03.98)			
(30) Priority Data: 60/040,623 7 March 1997 (07.03.97) US			
(71) Applicant: METABASIS THERAPEUTICS, INC. [US/US]; 9390 Towne Centre Drive, San Diego, CA 92121 (US).			
(72) Inventors: DANG, Qun; 7825 Roan Road, San Diego, CA 92129 (US). ERION, Mark, D.; 13455 Mango Drive, Del Mar, CA 92104 (US). REDDY, M., Rami; 9515 Easter Way #3, San Diego, CA 92121 (US). ROBINSON, Edward, D.; 3567 Tony Drive, San Diego, CA 92122 (US). KASIBHATLA, Srinivas, Rao; 10788 Glendover Lane, San Diego, CA 92126 (US). REDDY, K., Raja; 4146 Federman Lane, San Diego, CA 92130 (US).		Published <i>With international search report.</i>	
(74) Agents: WOLFF, Jessica, R. et al.; Lyon & Lyon LLP, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071-2066 (US).			
(54) Title: NOVEL PURINE INHIBITORS OF FRUCTOSE-1,6-BISPHOSPHATASE			
(57) Abstract			
<p>Novel purine compounds of structure (1) and their use as fructose-1,6-bisphosphatase inhibitors are described, wherein A is selected from the group consisting of -NR⁶₂, NHSO₂R³, -OR⁵, -SR⁵, halogen, lower alkyl, -CON(R⁴)₂, guanidine, amidine, -H, and perhaloalkyl; E is selected from the group consisting of -H, halogen, lower alkylthio, lower perhaloalkyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, -CN, and -NR⁷₂; X is selected from the group consisting of alkylamino, alkyl, alkenyl, alkynyl, alkyl(carboxyl), alkyl(hydroxy), alkyl(phosphonate), alkyl(sulfonate), aryl, alkylaminoalkyl, alkoxyalkyl, alkylthioalkyl, alkylthio, alicyclic, 1,1-dihaloalkyl, carbonylalkyl, aminocarbonylamino, alkylaminocarbonyl, alkylcarbonylamino, aralkyl, and alkylaryl, all optionally substituted; or together with Y forms a cyclic group including cyclic alkyl, heterocyclic, and aryl; Y is selected from the group consisting of -H, alkyl, alkenyl, alkynyl, aryl, alicyclic, aralkyl, aryloxyalkyl, alkoxyalkyl, -C(O)R³, -S(O)₂R³, -CO(O)-OR³, -CONHR³, -NR²₂, and -OR³, all except H are optionally substituted; or together with X forms a cyclic group including aryl, cyclic alkyl, and heterocyclic; and pharmaceutically acceptable prodrugs and salts thereof.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NOVEL PURINE INHIBITORS OF FRUCTOSE-1,6-BISPHOSPHATASE**Field of the Invention**

5 This invention relates to novel purine compounds that are inhibitors of Fructose-1,6-bisphosphatase at the AMP site. The invention also relates to the preparation and use of these purine analogs in the treatment of diabetes, and other diseases where the inhibition of gluconeogenesis, control of blood glucose levels, reduction in glycogen stores, or reduction in insulin levels is beneficial.

10

Background and Introduction to the Invention

Diabetes mellitus (or diabetes) is one of the most prevalent diseases in the world today. Diabetes patients have been divided into two classes, namely type I or insulin-dependent diabetes mellitus and type II or non-insulin dependent diabetes mellitus (NIDDM). Non-insulin-dependent diabetes mellitus (NIDDM) accounts for approximately 90% of all diabetics and is estimated to affect 12-14 million adults in the U. S. alone (6.6% of the population). NIDDM is characterized by both fasting hyperglycemia and exaggerated postprandial increases in plasma glucose levels. NIDDM is associated with a variety of long-term complications, including microvascular diseases such as retinopathy, nephropathy and neuropathy, and macrovascular diseases such as coronary heart disease. Numerous studies in animal models demonstrate a causal relationship between long term complications and hyperglycemia. Recent results from the Diabetes Control and Complications Trial (DCCT) and the Stockholm Prospective Study demonstrate this relationship for the first time in man by showing that insulin-dependent diabetics with tighter glycemic control are at substantially lower risk for development and progression of these complications. Tighter control is also expected to benefit NIDDM patients.

30 Current therapies used to treat NIDDM patients entail both controlling lifestyle risk factors and pharmaceutical intervention. First-line therapy for NIDDM is typically a tightly-controlled regimen of diet and exercise since an overwhelming number of NIDDM patients are overweight or obese ($\approx 67\%$) and since weight loss can improve insulin secretion, insulin sensitivity and lead to normoglycemia. Normalization of blood glucose occurs in less than 30% of these patients due to poor compliance and poor response. Patients with

35

hyperglycemia not controlled by diet alone are subsequently treated with oral hypoglycemics or insulin. Until recently, the sulfonylureas were the only class of oral hypoglycemic agents available for NIDDM. Treatment with sulfonylureas leads to effective blood glucose lowering in only 70% of patients and only 40% after 10 years of therapy. Patients that fail to respond to diet and sulfonylureas are subsequently treated with daily insulin injections to gain adequate glycemic control.

Although the sulfonylureas represent a major therapy for NIDDM patients, four factors limit their overall success. First, as mentioned above, a large segment of the NIDDM population do not respond adequately to sulfonylurea therapy (*i.e.* primary failures) or become resistant (*i.e.* secondary failures). This is particularly true in NIDDM patients with advanced NIDDM since these patients have severely impaired insulin secretion. Second, sulfonylurea therapy is associated with an increased risk of severe hypoglycemic episodes. Third, chronic hyperinsulinemia has been associated with increased cardiovascular disease although this relationship is considered controversial and unproven. Last, sulfonylureas are associated with weight gain, which leads to worsening of peripheral insulin sensitivity and thereby can accelerate the progression of the disease.

Recent results from the U.K. Diabetes prospective study also showed that patients undergoing maximal therapy of a sulfonylurea, metformin, or a combination of the two, were unable to maintain normal fasting glycemia over the six year period of the study. U.K. Prospective Diabetes Study 16. Diabetes, 44:1249-158 (1995). These results further illustrate the great need for alternative therapies. Three therapeutic strategies that could provide additional health benefits to NIDDM patients beyond the currently available therapies, include drugs that would: (i) prevent the onset of NIDDM; (ii) prevent diabetic complications by blocking detrimental events precipitated by chronic hyperglycemia; or (iii) normalize glucose levels or at least decrease glucose levels below the threshold reported for microvascular and macrovascular diseases.

Hyperglycemia in NIDDM is associated with two biochemical abnormalities, namely insulin resistance and impaired insulin secretion. The relative roles of these metabolic abnormalities in the pathogenesis of NIDDM has been the subject of numerous studies over the past several decades. Studies of offspring and siblings of NIDDM patients, mono- and dizygotic twins,

and ethnic populations with high incidence of NIDDM (e.g. Pima Indians) strongly support the inheritable nature of the disease.

Despite the presence of insulin resistance and impaired insulin secretion, fasting blood glucose (FBG) levels remain normal in pre-diabetic patients due to a state of compensatory hyperinsulinemia. Eventually, however, insulin secretion is inadequate and fasting hyperglycemia ensues. With time insulin levels decline. Progression of the disease is characterized by increasing FBG levels and declining insulin levels.

Numerous clinical studies have attempted to define the primary defect that accounts for the progressive increase in FBG. Results from these studies indicate that excessive hepatic glucose output (HGO) is the primary reason for the elevation in FBG with a significant correlation found for HGO and FBG once FBG exceeds 140 mg/dL. Kolterman, et al., J. Clin. Invest. 68:957, (1981); DeFronzo Diabetes 37:667 (1988).

HGO comprises glucose derived from breakdown of hepatic glycogen (glycogenolysis) and glucose synthesized from 3-carbon precursors (gluconeogenesis). A number of radioisotope studies and several studies using ¹³C-NMR spectroscopy have shown that gluconeogenesis contributes between 50-100% of the glucose produced by the liver in the postabsorptive state and that gluconeogenesis flux is excessive (2- to 3-fold) in NIDDM patients. Magnusson, et al. J. Clin. Invest. 90:1323-1327 (1992); Rothman, et al., Science 254: 573-76 (1991); Consoli, et al. Diabetes 38:550-557 (1989).

Gluconeogenesis from pyruvate is a highly regulated biosynthetic pathway requiring eleven enzymes (Figure 1). Seven enzymes catalyze reversible reactions and are common to both gluconeogenesis and glycolysis. Four enzymes catalyze reactions unique to gluconeogenesis, namely pyruvate carboxylase, phosphoenolpyruvate carboxykinase, fructose-1,6-bisphosphatase and glucose-6-phosphatase. Overall flux through the pathway is controlled by the specific activities of these enzymes, the enzymes that catalyzed the corresponding steps in the glycolytic direction, and by substrate availability. Dietary factors (glucose, fat) and hormones (insulin, glucagon, glucocorticoids, epinephrine) coordinatively regulate enzyme activities in the gluconeogenesis and glycolysis pathways through gene expression and post-translational mechanisms.

Of the four enzymes specific to gluconeogenesis, fructose-1,6-bisphosphatase (hereinafter "FBPase") is the most suitable target for a gluconeogenesis inhibitor based on efficacy and safety considerations. Studies indicate that nature uses the FBPase/PFK cycle as a major control point (metabolic switch) responsible for determining whether metabolic flux proceeds in the direction of glycolysis or gluconeogenesis. Claus, et al., Mechanisms of Insulin Action, Belfrage, P. editor, pp.305-321, Elsevier Science 1992; Regen, et al. J. Theor. Biol., 111:635-658 (1984); Pilkis, et al. Annu. Rev. Biochem., 57:755-783 (1988). FBPase is inhibited by fructose-2,6-bisphosphate in the cell. Fructose-2,6-bisphosphate binds to the substrate site of the enzyme. AMP binds to an allosteric site on the enzyme.

Synthetic inhibitors of FBPase have also been reported. McNiel reported that fructose-2,6-bisphosphate analogs inhibit FBPase by binding to the substrate site. J. Med. Chem., 106:7851 (1984); U.S. Patent No. 4,968,790 (1984). These compounds, however, were relatively weak and did not inhibit glucose production in hepatocytes presumably due to poor cell penetration.

Gruber reported that some nucleosides can lower blood glucose in the whole animal through inhibition of FBPase. These compounds exert their activity by first undergoing phosphorylation to the corresponding monophosphate. EP 0 427 799 B1.

Gruber et al. U.S. Patent No. 5,658,889 described the use of inhibitors of the AMP site of FBPase to treat diabetes.

European patent application EP 0 632 048 A1 discloses certain ethyl phosphonates of purine derivatives for use as antiviral and antineoplastic agents. These structures differ from the claimed compounds because they have no substitution on the C-8 of the purine. There is no suggestion that these compounds are inhibitors of FBPase.

Brief Description of the Drawings

FIG. 1 is a scheme depicting the eleven enzymes of the gluconeogenesis pathway.

FIG. 2 shows the dose-dependent inhibition of hPFBPase AMP and compounds 2.7, 2.13, and 2.5.

FIG. 3 shows the displacement of ³H-AMP from hPFBPase by ZMP and compound 2.2.

FIG. 4A depicts the reduction of glucose production from dihydroxyacetone in rat hepatocytes treated with compound 2.7.

FIG. 4B depicts the increase in the amount of fructose-1,6-bisphosphate in rat hepatocytes exposed to dihydroxyacetone and treated with compound 2.7.

FIG. 5 depicts a dose-dependent inhibition of glucose production in rat hepatocytes exposed to lactate and pyruvate by compounds 2.7 and 2.1.

FIG. 6 shows the effect of various doses of compound 16.4 on blood glucose levels in 18 hour fasted, normal rats.

FIG. 7 is a bar graph depicting the reduction in blood glucose levels in 18 hour fasted normal rats treated with compound 2.7 given at a dose of 20 mg/kg i.p.

FIG. 8 is a bar graph depicting the increased accumulation of fructose-1,6-bisphosphate in the liver of 18-hour fasted rats treated with compound 2.7.

FIG. 9A is a bar graph depicting a reduction in blood glucose levels in 24 hour fasted Zucker Diabetic Fatty rats treated with compound 2.7.

FIG. 9B is a bar graph depicting the percentage change in blood glucose levels in 24 hour fasted Zucker Diabetic Fatty rats treated with compound 2.7.

FIG. 10 depicts the inhibition of gluconeogenesis from ^{14}C -bicarbonate in 24 hour fasted Diabetic Fatty rats treated with compound 2.7.

FIG. 11A depicts the dose-dependent inhibition of glucose production in rat hepatocytes by compound 16.4, a prodrug of compound 2.7.

FIG. 11B shows the intracellular generation of compound 2.7 in rat hepatocytes treated with compound 16.4, a prodrug, to inhibit glucose production in rat hepatocytes.


Summary of the Invention

The present invention is directed towards novel purine compounds which bind the AMP site and are potent FBPase inhibitors. In another aspect, the present invention is directed to the preparation of these novel purine compounds and to the *in vitro* and *in vivo* FBPase inhibitory activity of these compounds. Another aspect of the present invention is directed to the clinical use of the novel FBPase inhibitors as a method of treatment or prevention of diseases responsive to inhibition of gluconeogenesis and in diseases responsive to lowered blood glucose levels.

Gruber et al. U.S. patent application Serial Number 08/355,836 (allowed) described the use of inhibitors of the AMP site of FBPase to treat diabetes.

The compounds are also useful in treating or preventing excess glycogen storage diseases and insulin dependent diseases such as cardiovascular diseases including atherosclerosis.

The invention comprises the novel purine analogs as specified below in formula 1. Also included in the scope of the present invention are prodrugs of the compounds of formula 1.



Formula 1

Since these compounds may have asymmetric centers, the present invention is directed not only to racemic mixtures of these compounds, but also to individual stereoisomers. The present invention also includes pharmaceutically acceptable and/or useful salts of the compounds of formula 1, including acid addition salts and basic salts. The present inventions also encompass prodrugs of compounds of formula 1.

Definitions

In accordance with the present invention and as used herein, the following terms are defined with the following meanings, unless explicitly stated otherwise.

The term "aryl" refers to aromatic groups which have at least one ring having a conjugated pi electron system and includes carbocyclic aryl, heterocyclic aryl and biaryl groups, all of which may be optionally substituted.

Carbocyclic aryl groups are groups wherein the ring atoms on the aromatic ring are carbon atoms. Carbocyclic aryl groups include monocyclic carbocyclic aryl groups and polycyclic or fused compounds such as optionally substituted naphthyl groups.

Heterocyclic aryl groups are groups having from 1 to 4 heteroatoms as ring atoms in the aromatic ring and the remainder of the ring atoms being carbon atoms. Suitable heteroatoms include oxygen, sulfur, and nitrogen. Suitable heteroaryl groups include furanyl, thienyl, pyridyl, pyrrolyl, N-lower alkyl pyrrolyl, pyridyl-N-oxide, pyrimidyl, pyrazinyl, imidazolyl, and the like, all optionally substituted.

The term "biaryl" represents aryl groups containing more than one aromatic ring including both fused ring systems and aryl groups substituted with other aryl groups.

The term "alicyclic" means compounds which combine the properties of aliphatic and cyclic compounds and include but are not limited to aromatic, cycloalkyl and bridged cycloalkyl compounds. The cyclic compound includes heterocycles. Cyclohexenylethyl, cyclohexanylethyl, and norbornyl are suitable alicyclic groups. Such groups may be optionally substituted.

The term "optionally substituted" or "substituted" includes groups substituted by one to four substituents, independently selected from lower alkyl, lower aryl, lower aralkyl, lower alicyclic, hydroxy, lower alkoxy, lower aryloxy, perhaloalkoxy, aralkoxy, heteroaryl, heteroaryloxy, heteroarylalkyl, heteroaralkoxy, azido, amino, guanidino, halogen, lower alkylthio, oxa, ketone, carboxy esters, carboxyl, carboxamido, nitro, acyloxy, alkylamino, aminoalkyl, alkylaminoaryl, alkylaryl, alkylaminoalkyl, alkoxyaryl, arylamino, aralkylamino, phosphonate, sulfonate, carboxamidoalkylaryl, carboxamidoaryl, hydroxyalkyl, haloalkyl, alkylaminoalkylcarboxy, aminocarboxamidoalkyl, cyano, lower alkoxyalkyl, and lower perhaloalkyl.

The term "aralkyl" refers to an alkyl group substituted with an aryl group. Suitable aralkyl groups include benzyl, picolyl, and the like, and may be optionally substituted.

The term "lower" referred to herein in connection with organic radicals or compounds respectively defines such as with up to and including 10, preferably up to and including 6, and advantageously one to four carbon atoms. Such groups may be straight chain, branched, or cyclic.

The terms "arylamino" (a), and "aralkylamino" (b), respectively, refer to the group -NRR' wherein respectively, (a) R is aryl and R' is hydrogen, alkyl, aralkyl or aryl, and (b) R is aralkyl and R' is hydrogen or aralkyl, aryl, alkyl.

The term "acyl" refers to -C(O)R where R is alkyl and aryl.

The term "carboxy esters" refers to $-C(O)OR$ where R is alkyl, aryl, aralkyl, and alicyclic, all optionally substituted.

The term "oxa" refers to $=O$ in an alkyl group.

The term "alkylamino" refers to $-NRR'$ where R and R' are independently
5 selected from hydrogen or alkyl.

The term "carbonylamine" or "carbonylamino" refers to $-CONR_2$ where each R is independently hydrogen or alkyl.

The term "halogen" or "halo" refers to $-F$, $-Cl$, $-Br$ and $-I$.

The term "oxyalkylamino" refers to $-O-alk-NR-$, where "alk" is an alkylene
10 group and R is H or alkyl.

The term "alkylsulfonate" refers to the group $-alk-S(O)_2-O-$ where "alk" is an alkylene group.

The term "alkylaminoalkylcarboxy" refers to the group $-alk-NR-alk-C(O)-O-$ where "alk" is an alkylene group, and R is a H or lower alkyl.

15 The term "alkylaminocarbonyl" refers to the group $-alk-NR-C(O)-$ where "alk" is an alkylene group, and R is a H or lower alkyl.

The term "oxyalkyl" refers to the group $-O-alk-$ where "alk" is an alkylene group.

The term "alkylcarboxyalkyl" refers to the group $-alk-C(O)-O-alkyl$ where
20 each alk is independently an alkylene group.

The term "alkyl" refers to saturated aliphatic groups including straight-chain, branched chain and cyclic groups. Alkyl groups may be optionally substituted.

The term "bidentate" refers to an alkyl group that is attached by its
25 terminal ends to the same atom to form a cyclic group. For example, propylene imine contains a bidentate propylene group.

The term "cyclic alkyl" refers to alkyl groups that are cyclic.

The term "heterocyclic" and "heterocyclic alkyl" refer to cyclic alkyl groups containing at least one heteroatom. Suitable heteroatoms include oxygen,
30 sulfur, and nitrogen. Heterocyclic groups may be attached through a heteroatom or through a carbon atom in the ring.

The term "alkenyl" refers to unsaturated groups which contain at least one carbon-carbon double bond and includes straight-chain, branched-chain and cyclic groups. Alkene groups may be optionally substituted.

The term "alkynyl" refers to unsaturated groups which contain at least one carbon-carbon triple bond and includes straight-chain, branched-chain and cyclic groups. Alkyne groups may be optionally substituted.

5 The term "alkylene" refers to a divalent straight chain, branched chain or cyclic saturated aliphatic radical.

The term "acyloxy" refers to the ester group $-O-C(O)R$, where R is H, alkyl, alkenyl, alkynyl, aryl, aralkyl, or alicyclic.

10 The term "alkylaryl" refers to the group -alk-aryl- where "alk" is an alkylene group. "Lower alkylaryl" refers to such groups where alkylene is lower alkyl.

The term "alkylamino" refers to the group -alk-NR- wherein "alk" is an alkylene group.

15 The term "alkyl(carboxyl)" refers to carboxyl substituted off the alkyl chain. Similarly, "alkyl(hydroxy)", "alkyl(phosphonate)", and "alkyl(sulfonate)" refers to substituents off the alkyl chain.

The term "alkylaminoalkyl" refers to the group -alk-NR-alk- wherein each "alk" is an independently selected alkylene, and R is H or lower alkyl. "Lower alkylaminoalkyl" refers to groups where each alkylene group is lower alkyl.

20 The term "alkylaminoaryl" refers to the group -alk-NR-aryl- wherein "alk" is an alkylene group. In "lower alkylaminoaryl", the alkylene group is lower alkyl.

The term "alkyloxyaryl" refers to an alkylene group substituted with an aryloxy group. In "lower alkyloxyaryl", the alkylene group is lower alkyl.

25 The term "alkylacylamino" refers to the group -alk-N-(COR)- wherein alk is alkylene and R is lower alkyl. In "lower alkylacylamino", the alkylene group is lower alkyl.

30 The term "alkoxyalkylaryl" refers to the group -alk-O-alk-aryl- wherein each "alk" is independently an alkylene group. "Lower alkoxyalkylaryl" refers to such groups where the alkylene group is lower alkyl.

The term "alkylacylaminoalkyl" refers to the group -alk-N-(COR)-alk- where each alk is an independently selected alkylene group. In "lower alkylacylaminoalkyl" the alkylene groups are lower alkyl.

35 The term "alkoxy" refers to the group -alk-O- wherein alk is an alkylene group.

The term "alkoxyalkyl" refers to the group -alk-O-alk- wherein each alk is an independently selected alkylene group. In "lower alkoxyalkyl", each alkylene is lower alkyl.

5 The term "alkylthio" refers to the group -alk-S- wherein alk is alkylene group.

The term "alkylthioalkyl" refers to the group -alk-S-alk- wherein each alk is an independently selected alkylene group. In "lower alkylthioalkyl" each alkylene is lower alkylene.

10 The term "aralkylamino" refers to an amine substituted with an aralkyl group.

The term "alkylcarboxamido" refers to the group -alk-C(O)N(R)- wherein alk is an alkylene group and R is H or lower alkyl.

15 The term "alkylcarboxamidoalkyl" refers to the group -alk-C(O)N(R)-alk- wherein each alk is an independently selected alkylene group and R is lower alkyl. In "lower alkylcarboxamidoalkyl" each alkylene is lower alkyl.

20 The term "alkylcarboxamidoalkylaryl" refers to the group -alk₁-C(O)-NH-alk₂Ar- wherein alk₁ and alk₂ are independently selected alkylene groups and alk₂ is substituted with an aryl group, Ar. In "lower alkylcarboxamidoalkylaryl", each alkylene is lower alkyl.

The term "heteroalicyclic" refers to an alicyclic group having 1 to 4 heteroatoms selected from nitrogen, sulfur, phosphorus and oxygen.

25 The term "aminocarboxamidoalkyl" refers to the group -NH-C(O)-N(R)-R wherein each R is an independently selected alkyl group. "Lower aminocarboxamidoalkyl" refers to such groups wherein each R is lower alkyl.

The term "heteroarylalkyl" refers to an alkyl group substituted with a heteroaryl group.

30 The term "perhalo" refers to groups wherein every C-H bond has been replaced with a C-halo bond on an aliphatic or aryl group. Suitable perhaloalkyl groups include -CF₃ and -CFCl₂.

The term "guanidine" refers to both -NR-C(NR)-NR₂ as well as -N=C(NR₂)₂ where each R group is independently selected from the group of -H, alkyl, alkenyl, alkynyl, aryl, and alicyclic, all optionally substituted.


The term "amidine" refers to $-C(NR)-NR_2$ where each R group is independently selected from the group of -H, alkyl, alkenyl, alkynyl, aryl, and alicyclic, all optionally substituted.

The term "pharmaceutically acceptable salt" includes salts of compounds of formula 1 and its prodrugs derived from the combination of a compound of this invention and an organic or inorganic acid or base.

The term "prodrug" as used herein refers to any compound that when administered to a biological system generates the "drug" substance either as a result of spontaneous chemical reaction(s) or by enzyme catalyzed or metabolic reaction(s). Reference is made to various prodrugs such as acyl esters, carbonates, and carbamates, included herein. The groups illustrated are exemplary, not exhaustive, and one skilled in the art could prepare other known varieties of prodrugs. Such prodrugs of the compounds of formula 1, fall within the scope of the present invention.

The term "prodrug ester" as employed herein includes, but is not limited to, the following groups and combinations of these groups:

[1] Acyloxyalkyl esters which are well described in the literature (Farquhar et al., J. Pharm. Sci. 72, 324-325 (1983)) and are represented by formula A




Formula A

25

wherein R, R', and R'' are independently H, alkyl, aryl, alkylaryl, and alicyclic; (see WO 90/08155; WO 90/10636).

[2] Other acyloxyalkyl esters are possible in which an alicyclic ring is formed such as shown in formula B. These esters have been shown to generate phosphorus-containing nucleotides inside cells through a postulated sequence of reactions beginning with deesterification and followed by a series of elimination reactions (e.g. Freed et al., Biochem. Pharm. 38: 3193-3198 (1989)).




Formula B

5

wherein R is -H, alkyl, aryl, alkylaryl, alkoxy, aryloxy, alkylthio, arylthio, alkylamino, arylamino, cycloalkyl, or alicyclic.

[3] Another class of these double esters known as
 10 alkyloxycarbonyloxymethyl esters, as shown in formula A, where R is alkoxy, aryloxy, alkylthio, arylthio, alkylamino, and arylamino; R', and R" are independently H, alkyl, aryl, alkylaryl, and alicyclic, have been studied in the area of β -lactam antibiotics (Tatsuo Nishimura et al. *J. Antibiotics*, **1987**, *40*(1), 81-90; for a review see Ferres, H., *Drugs of Today*, **1983**, *19*, 499.). More
 15 recently Cathy, M. S., et al. (Abstract from AAPS Western Regional Meeting, April, **1997**) showed that these alkyloxycarbonyloxymethyl ester prodrugs on (9-[(R)-2-phosphonomethoxy]propyl]adenine (PMPA) are bioavailable up to 30% in dogs.

20 [4] Aryl esters have also been used as phosphonate prodrugs (e.g. Erion, DeLambert et al., *J. Med. Chem.* 37: 498, 1994; Serafinowska et al., *J. Med. Chem.* 38: 1372, 1995). Phenyl as well as mono and poly-substituted phenyl phosphonate ester prodrugs have generated the parent phosphonic acid in studies conducted in animals and in man (Formula C). Another
 25 approach has been described where Y is a carboxylic ester ortho to the phosphate. Khamnei and Torrence, *J. Med. Chem.*; 39:4109-4115 (1996).




Formula C

wherein Y is H, alkyl, aryl, alkylaryl, alkoxy, acetoxy, halogen, amino, alkoxy carbonyl, hydroxy, cyano, alkylamino, and alicyclic.

5

[5] Benzyl esters have also been reported to generate the parent phosphonic acid. In some cases, using substituents at the para-position can accelerate the hydrolysis. Benzyl analogs with 4-acyloxy or 4-alkyloxy group [Formula D, X = H, OR or O(CO)R or O(CO)OR] can generate the 4-hydroxy compound more readily through the action of enzymes, *e.g.* oxidases, esterases, etc. Examples of this class of prodrugs are described by Mitchell et al., J. Chem. Soc. Perkin Trans. I 2345 (1992); Brook, et al. WO 91/19721.

10



15

Formula D

wherein X and Y are independently H, alkyl, aryl, alkylaryl, alkoxy, acetoxy, hydroxy, cyano, nitro, perhaloalkyl, halo, or alkyloxycarbonyl; and

20


R' and R'' are independently H, alkyl, aryl, alkylaryl, halogen, and alicyclic.

[6] Thio-containing phosphonate ester prodrugs have been described that are useful in the delivery of FBPase inhibitors to hepatocytes. These phosphonate ester prodrugs contain a protected thioethyl moiety as shown in formula E. One or more of the oxygens of the phosphonate can be esterified. Since the mechanism that results in de-esterification requires the generation of a free thiolate, a variety of thiol protecting groups are possible. For example, the disulfide is reduced by a reductase-mediated process (Puech et al., Antiviral Res., 22: 155-174 (1993)). Thioesters will also generate free thiolates after esterase-mediated hydrolysis. Benzaria, et al., J. Med. Chem., 39:4958 (1996). Cyclic analogs are also possible and were shown to liberate

25

30

phosphonate in isolated rat hepatocytes. The cyclic disulfide shown below has not been previously described and is novel.




Formula E

5

wherein Z is alkylcarbonyl, alkoxy carbonyl, arylcarbonyl, aryloxy carbonyl, or alkylthio.

Other examples of suitable prodrugs include proester classes exemplified by Biller and Magnin (U.S. Patent No. 5,157,027); Serafinowska et al. (J. Med. Chem. 38, 1372 (1995)); Starrett et al. (J. Med. Chem. 37, 1857 (1994)); Martin et al. J. Pharm. Sci. 76, 180 (1987); Alexander et al., Collect. Czech. Chem. Commun. 59, 1853 (1994)); and EPO patent application 0 632 048 A1. Some of the structural classes described are optionally substituted, including fused lactones attached at the omega position and optionally substituted 2-oxo-1,3-dioxolenes attached through a methylene to the phosphorus oxygen such as:



20

3-phthalidyl

2-oxotetrahydrofuran-5-yl


2-oxo-4,5-didehydro-1,3-dioxolanemethyl

wherein R is -H, alkyl, cycloalkyl, or alicyclic; and

25

wherein Y is -H, alkyl, aryl, alkylaryl, cyano, alkoxy, acetoxy, halogen, amino, alkylamino, alicyclic, and alkoxy carbonyl.

[7] Propyl phosphonate ester prodrugs can also be used to deliver
FBPase inhibitors into hepatocytes. These phosphonate ester prodrugs may
contain a hydroxyl and hydroxyl group derivatives at the 3-position of the propyl
group as shown in formula F. The R and X groups can form a cyclic ring system
5 as shown in formula F. One or more of the oxygens of the phosphonate can be
esterified.




10

Formula F

wherein R is alkyl, aryl, heteroaryl;
X is hydrogen, alkylcarbonyloxy, alkyloxycarbonyloxy; and
Y is alkyl, aryl, heteroaryl, alkoxy, alkylamino, alkylthio, halogen,
15 hydrogen, hydroxy, acetoxy, amino.

[8] The cyclic propyl phosphonate esters as in Formula G are shown
to activate to phosphonic acids. The activation of prodrug can be
mechanistically explained by *in vivo* oxidation and elimination steps. These
20 prodrugs inhibit glucose production in isolated rat hepatocytes and are also
shown to deliver FBPase inhibitors to the liver following oral administration.



25

Formula G

wherein

V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or

5 together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxycarboxy, or aryloxycarboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

10 together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxycarboxy, alkylthiocarboxy, hydroxymethyl, and aryloxycarboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;

Z is selected from the group consisting of $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOR}^3$, $-\text{CH}_2\text{OC(O)SR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{SR}^3$, $-\text{S(O)R}^3$, $-\text{CH}_2\text{N}_3$, $-\text{CH}_2\text{NR}^2_2$, $-\text{CH}_2\text{Ar}$,
15 $-\text{CH(Ar)OH}$, $-\text{CH(CH=CR}^2\text{R}^2)\text{OH}$, $-\text{CH(C}\equiv\text{CR}^2)\text{OH}$, and $-\text{R}^2$;

with the provisos that:

a) V, Z, W are not all $-\text{H}$; and

b) when Z is $-\text{R}^2$, then at least one of V and W is not $-\text{H}$ or $-\text{R}^9$;

R^2 is selected from the group consisting of R^3 and $-\text{H}$;

20 R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl; and

R^9 is selected from the group consisting of alkyl, aralkyl, and alicyclic.

25 [9] Phosphoramidate derivatives have been explored as potential phosphonate prodrugs (e.g. McGuigan et al., *Antiviral Res.* **1990**, *14*: 345; **1991**, *15*: 255. Serafinowska et al., *J. Med. Chem.*, **1995**, *38*, 1372). Most phosphoramidates are unstable under aqueous acidic conditions and are hydrolyzed to the corresponding phosphonic acids. Cyclic phosphoramidates
30 have also been studied as phosphonate prodrugs because of their potential for greater stability compared to non cyclic phosphoramidates (e.g. Starrett et al., *J. Med. Chem.*, **1994**, *37*: 1857).

Other prodrugs are possible based on literature reports such as
35 substituted ethyls for example, bis(trichloroethyl)esters as disclosed by McGuigan, et al. *Bioorg Med. Chem. Lett.*, 3:1207-1210 (1993), and the phenyl

and benzyl combined nucleotide esters reported by Meier, C. et al. Bioorg. Med. Chem. Lett., 7:99-104 (1997).

X group nomenclature as used herein in formula 1 describes the group attached to the phosphonate and ends with the group attached to the 2-position of the benzimidazole ring. For example, when X is alkylamino, the following structure is intended:




Y group nomenclature likewise ends with the group attached to the ring.

DETAILED DESCRIPTION OF THE INVENTION

Novel Purine Compounds

Preferred compounds of the present invention are inhibitors of the AMP site of FBPase of the following formula (1):



wherein

A is selected from the group consisting of $-\text{NR}^8_2$, NHSO_2R^3 , $-\text{OR}^5$, $-\text{SR}^5$, halogen, lower alkyl, $-\text{CON}(\text{R}^4)_2$, guanidine, amidine, $-\text{H}$, and perhaloalkyl;

E is selected from the group consisting of $-\text{H}$, halogen, lower alkylthio, lower perhaloalkyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, $-\text{CN}$, and $-\text{NR}^7_2$;

X is selected from the group consisting of alkylamino, alkyl, alkenyl, alkynyl, alkyl(carboxyl), alkyl(hydroxy), alkyl(phosphonate), alkyl(sulfonate), aryl, alkylaminoalkyl, alkoxyalkyl, alkylthioalkyl, alkylthio, alicyclic, 1,1-dihaloalkyl, carbonylalkyl, aminocarbonylamino, alkylaminocarbonyl,

alkylcarbonylamino, aralkyl, and alkylaryl, all optionally substituted; or together with Y forms a cyclic group including cyclic alkyl, heterocyclic, and aryl;


Y is selected from the group consisting of -H, alkyl, alkenyl, alkynyl, aryl, alicyclic, aralkyl, aryloxyalkyl, alkoxyalkyl, $-C(O)R^3$, $-S(O)_2R^3$, $-C(O)-OR^3$,

- 5 $-CONHR^3$, $-NR^2_2$, and $-OR^3$, all except H are optionally substituted; or together with X forms a cyclic group including aryl, cyclic alkyl, and heterocyclic;

R^1 is independently selected from the group consisting of

-H, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or thiocarbonate, $-C(R^2)_2$ -aryl, alkylaryl, $-C(R^2)_2OC(O)NR^2_2$,

- 10 $-NR^2-C(O)-R^3$, $-C(R^2)_2-OC(O)R^3$, $C(R^2)_2-O-C(O)OR^3$, $-C(R^2)_2OC(O)SR^3$, alkyl-S- $C(O)R^3$, alkyl-S-S-alkylhydroxy, and alkyl-S-S-S-alkylhydroxy, or together R^1 and R^1 are -alkyl-S-S-alkyl to form a cyclic group, or together R^1 and R^1 are



15

wherein

V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^3$; or

- 20 together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxy, alkoxy, or aryloxy, or aryloxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

- 25 together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxy, alkoxy, alkythiocarboxy, hydroxymethyl, and aryloxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;

- 30 Z is selected from the group consisting of $-CH_2OH$, $-CH_2OCOR^3$, $-CH_2OC(O)SR^3$, $-CH_2OCO_2R^3$, $-SR^3$, $-S(O)R^3$, $-CH_2N_3$, $-CH_2NR^2_2$, $-CH_2Ar$, $-CH(Ar)OH$, $-CH(CH=CR^2R^2)OH$, $-CH(C\equiv CR^2)OH$, and $-R^2$;

with the provisos that:

a) V, Z, W are not all -H; and

b) when Z is -R², then at least one of V and W is not -H or -R⁹;

R² is selected from the group consisting of R³ and -H;

5 R³ is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl;

R⁴ is independently selected from the group consisting of -H, lower alkyl, lower alicyclic, lower aralkyl, and lower aryl;

10 R⁵ is selected from the group consisting of lower alkyl, lower aryl, lower aralkyl, and lower alicyclic;

R⁶ is independently selected from the group consisting of -H, and lower alkyl;

R⁷ is independently selected from the group consisting of -H, lower alkyl, lower alicyclic, lower aralkyl, lower aryl, and -C(O)R¹⁰;

15 R⁸ is independently selected from the group consisting of -H, lower alkyl, lower aralkyl, lower aryl, lower alicyclic, -C(O)R¹⁰, or together they form a bidentate alkyl;

R⁹ is selected from the group consisting of alkyl, aralkyl, and alicyclic;

20 R¹⁰ is selected from the group consisting of -H, lower alkyl, -NH₂, lower aryl, and lower perhaloalkyl;

R¹¹ is selected from the group consisting of alkyl, aryl, -OH, -NH₂ and -OR³; and

pharmaceutically acceptable prodrugs and salts thereof.

25

Preferred Compounds of Formula 1

Suitable alkyl groups include groups having from 1 to about 20 carbon atoms. Suitable aryl groups include groups having from 1 to about 20 carbon atoms. Suitable aralkyl groups include groups having from 2 to about 21 carbon atoms. Suitable acyloxy groups include groups having from 1 to about 20 carbon atoms. Suitable alkylene groups include groups having from 1 to about 20 carbon atoms. Suitable alicyclic groups include groups having 3 to about 20 carbon atoms. Suitable heteroaryl groups include groups having from 1 to about 20 carbon atoms and from 1 to 5 heteroatoms, preferably
35 independently selected from nitrogen, oxygen, phosphorous, and sulfur. Suitable heteroalicyclic groups include groups having from 2 to about twenty

carbon atoms and from 1 to 5 heteroatoms, preferably independently selected from nitrogen, oxygen, phosphorous, and sulfur.

Preferred A groups include $-NR^8$, lower alkyl, lower perhaloalkyl, lower alkoxy, and halogen. Particularly preferred are $-NR^8$, and halogen. Especially preferred is $-NR^8$. Most preferred is $-NH_2$.


Preferred E groups include -H, halogen, lower perhaloalkyl, -CN, lower alkyl, lower alkoxy, and lower alkylthio. Particularly preferred E groups include -H, -SMe, -Et, and -Cl. Especially preferred is -H and $-SCH_3$.

Preferred X groups include alkylamino, alkyl, alkynyl, alkoxyalkyl, alkylthio, aryl, 1,1-dihaloalkyl, carbonylalkyl, heteroaryl, alkylcarbonylamino, and alkylaminocarbonyl. Particularly preferred is alkyl substituted with 1 to 3 substituents selected from halogen, phosphonate, $-CO_2H$, $-SO_3H$, and -OH. Particularly preferred are alkylaminocarbonyl, alkoxyalkyl, and heteroaryl. Preferred alkoxyalkyl groups include methoxymethyl. Preferred heteroaryl groups include furanyl, optionally substituted.

Preferred Y groups include aralkyl, alicyclic, alkyl, and aryl, all optionally substituted. Particularly preferred is lower alkyl. Particularly preferred Y groups include (2-naphthyl)methyl, cyclohexylethyl, phenylethyl, nonyl, cyclohexylpropyl, ethyl, cyclopropylmethyl, cyclobutylmethylphenyl, (2-methyl)propyl, neopentyl, cyclopropyl, cyclopentyl, (1-imidazolyl)propyl, 2-ethoxybenzyl, 1-hydroxy-2,2-dimethylpropyl, 1-chloro-2,2-dimethylpropyl, 2,2-dimethylbutyl, 2-(spiro-3,3'-dimethylcyclohex-4-enyl)propyl, and 1-methylnopentyl. Especially preferred is neopentyl and isobutyl.

Preferred R^4 and R^7 groups are -H, and lower alkyl. Particularly preferred are -H, and methyl.

Preferred R^1 groups include -H, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or thiocarbonate, $-C(R^2)_2$ -aryl, alkylaryl, $-C(R^2)_2OC(O)NR^2$, $-NR^2-C(O)-R^3$, $-C(R^2)_2-OC(O)R^3$, $C(R^2)_2-O-C(O)OR^3$, $-C(R^2)_2OC(O)SR^3$, alkyl-S-C(O) R^3 , alkyl-S-S-alkylhydroxy, and alkyl-S-S-S-alkylhydroxy, or together R^1 and R^1 are -alkyl-S-S-alkyl to form a cyclic group, or together R^1 and R^1 are



wherein

V and W are independently selected from the group consisting of
 5 hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or

together V and Z are connected to form a cyclic group containing 3-5
 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy,
 alkoxycarboxy, or aryloxycarboxy attached to a carbon atom that is three atoms
 10 from an oxygen attached to the phosphorus; or

together V and W are connected to form a cyclic group containing 3
 carbon atoms substituted with hydroxy, acyloxy, alkoxycarboxy,
 alkylthiocarboxy, hydroxymethyl, and aryloxycarboxy attached to a carbon atom
 that is three atoms from an oxygen attached to the phosphorus;

15 Z is selected from the group consisting of $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOR}^3$,
 $-\text{CH}_2\text{OC}(\text{O})\text{SR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{SR}^3$, $-\text{S}(\text{O})\text{R}^3$, $-\text{CH}_2\text{N}_3$, $-\text{CH}_2\text{NR}^2_2$, $-\text{CH}_2\text{Ar}$,
 $-\text{CH}(\text{Ar})\text{OH}$, $-\text{CH}(\text{CH}=\text{CR}^2\text{R}^2)\text{OH}$, $-\text{CH}(\text{C}\equiv\text{CR}^2)\text{OH}$, and $-R^2$;


with the provisos that:

- a) V, Z, W are not all $-\text{H}$; and
- 20 b) when Z is $-R^2$, then at least one of V and W is not $-\text{H}$ or $-R^9$;
 R^2 is selected from the group consisting of R^3 and $-\text{H}$;
 R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and
 aralkyl; and

R^9 is selected from the group consisting of alkyl, aralkyl, and
 25 alicyclic.

Preferred R^1 groups include $-\text{H}$, alkylaryl, aryl, $-\text{C}(\text{R}^2)_2\text{-aryl}$, and $-\text{C}(\text{R}^2)_2\text{-OC}(\text{O})\text{R}^3$. Preferred such R^1 groups include optionally substituted phenyl, optionally substituted benzyl, $-\text{H}$, $-\text{C}(\text{R}^2)_2\text{OC}(\text{O})\text{OR}^3$, and $-\text{C}(\text{R}^2)_2\text{OC}(\text{O})\text{R}^3$. Preferably, said alkyl groups are greater than 4 carbon atoms. Another
 30 preferred aspect is where at least one R^1 is aryl or $-\text{C}(\text{R}^2)_2\text{-aryl}$. Also particularly preferred are compounds where R^1 is alicyclic where the cyclic moiety contains

- carbonat or thiocarbonate. Another preferred aspect is when at least one R^1 is $-C(R^2)_2-OC(O)R^3$, $-C(R^2)_2-OC(O)OR^3$ or $-C(R^2)_2-OC(O)SR^3$. Also particularly preferred is when R^1 and R^1 together are optionally substituted, including fused, lactone attached at the omega position or are optionally substituted 2-oxo-1,3-dioxolenes attached through a methylene to the phosphorus oxygen. Also preferred is when at least one R^1 is -alkyl-S-S-alkylhydroxyl, -alkyl-S-C(O) R^3 , and -alkyl-S-S-S-alkylhydroxy, or together R^1 and R^1 are -alkyl-S-S-alkyl- to form a cyclic group. Also preferred is where R^1 and R^1 together are



10

to form a cyclic group,
wherein

- V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or
- together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxycarboxy, or aryloxycarboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or
- together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxycarboxy, alkylthiocarboxy, hydroxymethyl, and aryloxycarboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;
- Z is selected from the group consisting of $-CH_2OH$, $-CH_2OCOR^3$, $-CH_2OC(O)SR^3$, $-CH_2OCO_2R^3$, $-SR^3$, $-S(O)R^3$, $-CH_2N_3$, $-CH_2NR^2_2$, $-CH_2Ar$, $-CH(Ar)OH$, $-CH(CH=CR^2R^2)OH$, $-CH(C\equiv CR^2)OH$, and $-R^2$;
- with the provisos that:
- V, Z, W are not all -H; and
 - when Z is $-R^2$, then at least one of V and W is not -H or $-R^9$;
- R^2 is selected from the group consisting of R^3 and -H;

30

R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl; and


R^9 is selected from the group consisting of alkyl, aralkyl, and alicyclic.

- 5 Particularly preferred are such groups wherein V and W both form a 6-membered carbocyclic ring substituted with 0-4 groups, selected from the group consisting of hydroxy, acyloxy, alkoxycarbonyl, and alkoxy; and Z is R^2 . Also particularly preferred are such groups wherein V and W are hydrogen; and Z is selected from the group consisting of hydroxyalkyl, acyloxyalkyl, alkyloxyalkyl, 10 and alkoxycarboxyalkyl. Also particularly preferred are such groups wherein V and W are independently selected from the group consisting of hydrogen, optionally substituted aryl, and optionally substituted heteroaryl, with the proviso that at least one of V and W is optionally substituted aryl or optionally substituted heteroaryl.

15

In one preferred aspect, R^1 is not lower alkyl of 1-4 carbon atoms.

- In another preferred aspect, A is $-NR^8_2$ or halogen, E is -H, halogen, -CN, lower alkyl, lower perhaloalkyl, lower alkoxy, or lower alkylthio, X is alkylamino, alkyl, alkoxyalkyl, alkynyl, 1,1-dihaloalkyl, carbonylalkyl, alkyl(OH), 20 alkyl(sulfonate), alkylcarbonylamino, alkylaminocarbonyl, alkylthio, aryl, or heteroaryl, and R^4 and R^7 is -H or lower alkyl. Particularly preferred are such compounds where Y is aralkyl, aryl, alicyclic, or alkyl. Especially preferred are such compounds where R^1 and R^1 together are



25

wherein

V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or

- 30 together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy,

alkoxycarboxy, or aryloxy-carboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

- together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxycarboxy, alkylthiocarboxy, hydroxymethyl, and aryloxy-carboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;

Z is selected from the group consisting of $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOR}^3$, $-\text{CH}_2\text{OC(O)SR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{SR}^3$, $-\text{S(O)R}^3$, $-\text{CH}_2\text{N}_3$, $-\text{CH}_2\text{NR}^2_2$, $-\text{CH}_2\text{Ar}$, $-\text{CH(Ar)OH}$, $-\text{CH(CH=CR}^2\text{R}^2\text{)OH}$, $-\text{CH(C}\equiv\text{CR}^2\text{)OH}$, and $-\text{R}^2$;

- with the provisos that:

- a) V, Z, W are not all -H; and
- b) when Z is $-\text{R}^2$, then at least one of V and W is not -H or $-\text{R}^9$;
 R^2 is selected from the group consisting of R^3 and -H;
 R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and

- aralkyl; and

R^9 is selected from the group consisting of alkyl, aralkyl, and alicyclic.

- In another preferred aspect, A is $-\text{NR}^8_2$, E is -H, Cl-, or methylthio, and X is optionally substituted furanyl, or alkoxyalkyl. Particularly preferred are such compounds where A is $-\text{NH}_2$, X is 2,5-furanyl, or methoxymethyl, and Y is lower alkyl. Most preferred are such compounds where E is H, X is 2,5-furanyl, and Y is neopentyl; those where E is $-\text{SCH}_3$, X is 2,5-furanyl, and Y is isobutyl; and those where E is -H, X is 2,5-furanyl, and Y is 1-(3-chloro-2,2-dimethyl)-propyl. Especially preferred are such compounds where R^1 is $-\text{CH}_2\text{O-C(O)-C(CH}_3\text{)}_3$.

25

In the following examples of preferred compounds, the following prodrugs are preferred:

- Acyloxyalkyl esters;
- Alkoxycarbonyloxyalkyl esters;
- 30 Aryl esters;
- Benzyl and substituted benzyl esters;
- Disulfide containing esters;
- Substituted (1,3-dioxolen-2-one)methyl esters;
- Substituted 3-phthalidyl esters;
- 35 Cyclic-[2'-hydroxymethyl]-1,3-propanyl diesters and hydroxy protected forms;

Lactone type esters; and all mixed esters resulted from possible combinations of above esters.

- Bis-pivaloyloxymethyl esters;
- 5 Bis-isobutyryloxymethyl esters;
- Cyclic-[2'-hydroxymethyl]-1,3-propanyl diester;
- Cyclic-[2'-acetoxymethyl]-1,3-propanyl diester;
- Cyclic-[2'-methyloxycarbonyloxymethyl]-1,3-propanyl diester;
- Bis-benzoylthiomethyl esters;
- 10 Bis-benzoylthioethyl esters;
- Bis-benzoyloxymethyl esters;
- Bis-*p*-fluorobenzoyloxymethyl esters;
- Bis-6-chloronicotinoyloxymethyl esters;
- Bis-5-bromonicotinoyloxymethyl esters;
- 15 Bis-thiophenecarbonyloxymethyl esters;
- Bis-2-furoyloxymethyl esters;
- Bis-3-furoyloxymethyl esters;
- Diphenyl esters;
- Bis-(4-methoxyphenyl) esters;
- 20 Bis-(2-methoxyphenyl) esters;
- Bis-(2-ethoxyphenyl) esters;
- Mono-(2-ethoxyphenyl) esters;
- Bis-(4-acetamidophenyl) esters;
- Bis-(4-aceyloxyphenyl) esters;
- 25 Bis-(4-hydroxyphenyl) esters;
- Bis-(2-acetoxyphenyl) esters;
- Bis-(3-acetoxyphenyl) esters;
- Bis-(4-morpholinophenyl) esters;
- Bis-[4-(1-triazolophenyl) esters;
- 30 Bis-(3-*N,N*-dimethylaminophenyl) esters;
- Bis-(2-tetrahydronaphthyl) esters;
- Bis-(3-chloro-4-methoxy)benzyl esters;
- Bis-(3-bromo-4-methoxy)benzyl esters;
- Bis-(3-cyano-4-methoxy)benzyl esters;
- 35 Bis-(3-chloro-4-acetoxy)benzyl esters;
- Bis-(3-bromo-4-acetoxy)benzyl esters;

- Bis-(3-cyano-4-acetoxy)benzyl esters;
Bis-(4-chloro)benzyl esters;
Bis-(4-acetoxy)benzyl esters;
Bis-(3,5-dimethoxy-4-acetoxy)benzyl esters;
5 Bis-(3-methyl-4-acetoxy)benzyl esters;
Bis-(benzyl)esters;
Bis-(3-methoxy-4-acetoxy)benzyl esters;
Bis-(3-chloro-4-acetoxy)benzyl esters;
cyclic-(2,2-dimethylpropyl)phosphonoamidate;
10 cyclic-(2-hydroxymethylpropyl) ester;
Bis-(6'-hydroxy-3',4'-disulfide)hexyl esters;
Bis-(6'-acetoxy-3',4'-disulfide)hexyl esters;
(3',4'-Dithia)cyclononane esters;
Bis-(5-methyl-1,3-dioxolen-2-one-4-yl)methyl esters;
15 Bis-(5-ethyl-1,3-dioxolen-2-one-4-yl)methyl esters;
Bis-(5-tert-butyl-1,3-dioxolen-2-one-4-yl)methyl esters;
Bis-3-(5,6,7-trimethoxy)phthalidyl esters;
Bis-(cyclohexyloxycarbonyloxymethyl) esters;
Bis-(isopropylloxycarbonyloxymethyl) esters;
20 Bis-(ethylloxycarbonyloxymethyl) esters;
Bis-(methyloxycarbonyloxymethyl) esters;
Bis-(isopropylthiocarbonyloxymethyl) esters;
Bis-(phenyloxycarbonyloxymethyl) esters;
Bis-(benzyloxycarbonyloxymethyl) esters;
25 Bis-(phenylthiocarbonyloxymethyl) esters;
Bis-(*p*-methoxyphenyloxycarbonyloxymethyl) esters;
Bis-(*m*-methoxyphenyloxycarbonyloxymethyl) esters;
Bis-(*o*-methoxyphenyloxycarbonyloxymethyl) esters;
Bis-(*o*-methylphenyloxycarbonyloxymethyl) esters;
30 Bis-(*p*-chlorophenyloxycarbonyloxymethyl) esters;
Bis-(1,4-biphenyloxycarbonyloxymethyl) esters;
Bis-[(2-phthalimidoethyl)oxycarbonyloxymethyl]esters;
Bis-(*N*-Phenyl, *N*-methylcarbamoxyloxymethyl) esters;
Bis-(2-trichloroethyl) esters;
35 Bis-(2-bromoethyl) esters;
Bis-(2-iodoethyl) esters;


- Bis-(2-azidoethyl) esters;
- Bis-(2-acetoxyethyl) esters;
- Bis-(2-aminoethyl) esters;
- Bis-(2-*N,N*-diaminoethyl) esters;
- 5 Bis-(2-aminoethyl) esters;
- Bis-(methoxycarbonylmethyl) esters;
- Bis-(2-aminoethyl) esters;
- Bis- [*N,N*-di(2-hydroxyethyl)]amidomethylesters;
- Bis-(2-aminoethyl) esters;
- 10 Bis-(2-methyl-5-thiozolomethyl) esters;
- Bis-(bis-2-hydroxyethylamidomethyl) esters.

Most preferred are the following:

- 15 Bis-pivaloyloxymethyl esters;
- Bis-isobutyryloxymethyl esters;
- cyclic-(2-hydroxymethylpropyl) ester;
- cyclic-(2-acetoxymethylpropyl) ester;
- cyclic-(2-methyloxycarbonyloxymethylpropyl) ester;
- 20 cyclic-(2-cyclohexylcarbonyloxymethylpropyl)ester;
- cyclic-(2-aminomethylpropyl)ester;
- cyclic-(2-azidomethylpropyl)ester;
- Bis-benzoylthiomethyl esters;
- Bis-benzoylthioethylesters;
- 25 Bis-benzoyloxymethyl esters;
- Bis-*p*-fluorobenzoyloxymethyl esters;
- Bis-6-chloronicotinoyloxymethyl esters;
- Bis-5-bromonicotinoyloxymethyl esters;
- Bis-thiophenecarbonyloxymethyl esters;
- 30 Bis-2-furoyloxymethyl esters;
- Bis-3-furoyloxymethyl esters;
- Diphenyl esters;
- Bis-(2-methyl)phenyl esters;
- Bis-(2-methoxy)phenyl esters;
- 35 Bis-(2-ethoxy)phenyl esters;
- Bis-(4-methoxy)phenyl esters;

- Bis-(3-bromo-4-methoxy)benzyl esters;
Bis-(4-acetoxy)benzyl esters;
Bis-(3,5-dimethoxy-4-acetoxy)benzyl esters;
Bis-(3-methyl-4-acetoxy)benzyl esters;
5 Bis-(3-methoxy-4-acetoxy)benzyl esters;
Bis-(3-chloro-4-acetoxy)benzyl esters;
Bis-(cyclohexyloxycarbonyloxymethyl) esters;
Bis-(isopropylloxycarbonyloxymethyl) esters;
Bis-(ethyloxycarbonyloxymethyl) esters;
10 Bis-(methyloxycarbonyloxymethyl) esters;
Bis-(isopropylthiocarbonyloxymethyl) esters;
Bis-(phenyloxycarbonyloxymethyl) esters;
Bis-(benzyloxycarbonyloxymethyl) esters;
Bis-(phenylthiocarbonyloxymethyl) esters;
15 Bis-(*p*-methoxyphenyloxycarbonyloxymethyl) esters;
Bis-(*m*-methoxyphenyloxycarbonyloxymethyl) esters;
Bis-(*o*-methoxyphenyloxycarbonyloxymethyl) esters;
Bis-(*o*-methylphenyloxycarbonyloxymethyl) esters;
Bis-(*p*-chlorophenyloxycarbonyloxymethyl) esters;
20 Bis-(1,4-biphenyloxycarbonyloxymethyl) esters;
Bis-[(2-phthalimidoethyl)oxycarbonyloxymethyl]esters;
Bis-(6'-hydroxy-3',4'-disulfide)hexyl esters; and
(3',4'-Disulfide)cyclononane esters.
Bis-(2-bromoethyl) esters;
25 Bis-(2-aminoethyl) esters;
Bis-(2-*N,N*-diaminoethyl) esters;

Examples of preferred compounds include, but are not limited to those described in Table 1 including salts and prodrugs thereof:

Table Compound No.	Synthetic Example No.				
		A	E	Y	X
1	2.7	NH ₂	H	neopentyl	2,5-furanyl
2		H	H	neopentyl	2,5-furanyl
3		Me	H	neopentyl	2,5-furanyl
4		Et	H	neopentyl	2,5-furanyl
5		Pr	H	neopentyl	2,5-furanyl
6		I	H	neopentyl	2,5-furanyl
7		Br	H	neopentyl	2,5-furanyl
8		Cl	H	neopentyl	2,5-furanyl
9		F	H	neopentyl	2,5-furanyl
10		OMe	H	neopentyl	2,5-furanyl
11		OEt	H	neopentyl	2,5-furanyl
12		OPr	H	neopentyl	2,5-furanyl
13		SMe	H	neopentyl	2,5-furanyl
14		SEt	H	neopentyl	2,5-furanyl
15		SPr	H	neopentyl	2,5-furanyl
16		SBn	H	neopentyl	2,5-furanyl
17		OBn	H	neopentyl	2,5-furanyl
18		NHMe	H	neopentyl	2,5-furanyl
19		NHEt	H	neopentyl	2,5-furanyl
20		NH-cPr	H	neopentyl	2,5-furanyl
21		NHOH	H	neopentyl	2,5-furanyl
22		NHNH ₂	H	neopentyl	2,5-furanyl
23		NHCHO	H	neopentyl	2,5-furanyl
24		NHAc	H	neopentyl	2,5-furanyl
25		NHCOCF ₃	H	neopentyl	2,5-furanyl
26		NHSO ₂ Me	H	neopentyl	2,5-furanyl
27		CONH ₂	H	neopentyl	2,5-furanyl
28		ONH ₂	H	neopentyl	2,5-furanyl
29	22.2	NH ₂	SMe	neopentyl	2,5-furanyl
30		H	SMe	neopentyl	2,5-furanyl
31		Me	SMe	neopentyl	2,5-furanyl
32		Et	SMe	neopentyl	2,5-furanyl
33		Pr	SMe	neopentyl	2,5-furanyl
34		I	SMe	neopentyl	2,5-furanyl
35		Br	SMe	neopentyl	2,5-furanyl
36		Cl	SMe	neopentyl	2,5-furanyl
37		F	SMe	neopentyl	2,5-furanyl
38		OMe	SMe	neopentyl	2,5-furanyl
39		OEt	SMe	neopentyl	2,5-furanyl
40		OPr	SMe	neopentyl	2,5-furanyl

41		SMe	SMe	ne pentyl	2,5-furanyl
42		SEt	SMe	neopentyl	2,5-furanyl
43		SPr	SMe	neopentyl	2,5-furanyl
44		SBn	SMe	neopentyl	2,5-furanyl
45		OBn	SMe	neopentyl	2,5-furanyl
46		NHMe	SMe	neopentyl	2,5-furanyl
47		NHEt	SMe	neopentyl	2,5-furanyl
48		NH-cPr	SMe	neopentyl	2,5-furanyl
49		NHOH	SMe	neopentyl	2,5-furanyl
50		NHNH2	SMe	neopentyl	2,5-furanyl
51		NHCHO	SMe	neopentyl	2,5-furanyl
52		NHAc	SMe	neopentyl	2,5-furanyl
53		NHCOCF3	SMe	neopentyl	2,5-furanyl
54		NHSO2Me	SMe	neopentyl	2,5-furanyl
55		CONH2	SMe	neopentyl	2,5-furanyl
56		ONH2	SMe	neopentyl	2,5-furanyl
57		NH2	H	isobutyl	2,5-furanyl
58	2.6	NH2	H	isopropyl	2,5-furanyl
59	2.5	NH2	H	ethyl	2,5-furanyl
60		NH2	H	methyl	2,5-furanyl
61	2.9	NH2	H	cyclopropyl	2,5-furanyl
62		NH2	H	cyclobutyl	2,5-furanyl
63	2.10	NH2	H	cyclopentyl	2,5-furanyl
64		NH2	H	cyclohexyl	2,5-furanyl
65		NH2	H	cycloheptanyl	2,5-furanyl
66		NH2	H	cyclopropylmethyl	2,5-furanyl
67		NH2	H	cyclobutylmethyl	2,5-furanyl
68		NH2	H	cyclopentylmethyl	2,5-furanyl
69		NH2	H	2-cyclopropylethyl	2,5-furanyl
70		NH2	H	2-cyclobutylethyl	2,5-furanyl
71		NH2	H	2-cyclopentylethyl	2,5-furanyl
72	2.2	NH2	H	2-cyclohexylethyl	2,5-furanyl
73	2.1	NH2	H	2-phenylethyl	2,5-furanyl
74		NH2	H	benzyl	2,5-furanyl
75		NH2	H	phenyl	2,5-furanyl
76		NH2	H	D-ribosyl	2,5-furanyl
77		NH2	H	H	2,5-furanyl
78		NH2	H	1-naphthylmethyl	2,5-furanyl
79	2.3	NH2	H	2-naphthylmethyl	2,5-furanyl
80		NH2	H	3-cyclopropylpropyl	2,5-furanyl
81		NH2	H	3-cyclobutylpropyl	2,5-furanyl
82		NH2	H	3-cyclopentylpropyl	2,5-furanyl
83		NH2	H	3-cyclohexylpropyl	2,5-furanyl
84	2.4	NH2	H	2,2-diphenylethyl	2,5-furanyl
85	2.8	NH2	H	adamentylmethyl	2,5-furanyl
86	2.11	NH2	H	2-ethoxybenzyl	2,5-furanyl
87	2.13	NH2	H	2,2-dimethyl-3-hydroxy-1-propyl	2,5-furanyl

88	2.12	NH2	H	2,2-dimethyl-3-dimethylamino-1-propyl	2,5-furanyl
89	2.14	NH2	H	2,2-dimethyl-3-chloro-1-propyl	2,5-furanyl
90	2.15	NH2	H	3,3-dimethyl-1-butyl	2,5-furanyl
91	2.17	NH2	H	1,2,2-trimethyl-1-propyl	2,5-furanyl
92	2.16	NH2	H	1,5,5-trimethyl-3-ene-1-cyclohexylmethyl	2,5-furanyl
93		NH2	H	4-pyrimidylmethyl	2,5-furanyl
94		NH2	H	2-(4-pyrimidyl)ethyl	2,5-furanyl
95		NH2	H	5-pyrimidylmethyl	2,5-furanyl
96		NH2	H	2-(5-pyrimidyl)ethyl	2,5-furanyl
97		NH2	H	2-pyrimidylmethyl	2,5-furanyl
98		NH2	H	2-(2-pyrimidyl)ethyl	2,5-furanyl
99		NH2	H	2-pyridylmethyl	2,5-furanyl
100		NH2	H	2-(2-pyridyl)ethyl	2,5-furanyl
101		NH2	H	3-pyridylmethyl	2,5-furanyl
102		NH2	H	2-(3-pyridyl)ethyl	2,5-furanyl
103		NH2	H	4-pyridylmethyl	2,5-furanyl
104		NH2	H	2-(4-pyridyl)ethyl	2,5-furanyl
105		NH2	H	2-carbamoyl ethyl	2,5-furanyl
106		NH2	H	1-(2-carbamoyl)propyl	2,5-furanyl
107		NH2	H	neopentyl	CONHCH2
108		NH2	H	neopentyl	CONHCH2CH2
109		NH2	H	neopentyl	CH2CH2CH2
110		NH2	H	neopentyl	CH2CH2CF2
111	5.5	NH2	H	neopentyl	CH2OCH2
112		NH2	H	neopentyl	CH2OCF2
113		NH2	H	neopentyl	CF2CF2CF2
114		NH2	H	neopentyl	acetylene
115		NH2	H	neopentyl	SCH2
116		NH2	H	neopentyl	SCH2CH2
117		NH2	H	neopentyl	CH2SCH2
118		NH2	H	neopentyl	NHCH2CH2
119		NH2	H	neopentyl	N(Ac)CH2CH2
120		NH2	H	neopentyl	N(Bz)CH2CH2
121		NH2	H	neopentyl	N(Me)CH2CH2
122		NH2	H	neopentyl	N(Bn)CH2CH2
123		NH2	H	neopentyl	NHCOCH2
124		NH2	H	neopentyl	NHCOCH2CH2
125		NH2	H	neopentyl	NHCOCF2
126		NH2	H	neopentyl	NHSO2CH2
127		NH2	H	neopentyl	N(Me)COCH2
128		NH2	H	neopentyl	N(Bn)COCH2
129		NH2	H	neopentyl	NHOCH2
130		NH2	H	neopentyl	CH2CH2CH(OH)
131		NH2	H	neopentyl	CH2CH2CH(CO2H)

132		NH ₂	H	n opentyl	CH ₂ CH ₂ CH(SO ₃ H)
133		NH ₂	H	neopentyl	CH ₂ CH ₂ CH(PO ₃ H ₂)
134	20.1	NH ₂	H	neopentyl	CH ₂ -(1,2-imidazolyl)
135		NH ₂	H	neopentyl	CH ₂ -(1,2-pyrrolyl)
136		NH ₂	H	neopentyl	CSNHCH ₂
137		NH ₂	H	neopentyl	2,5-tetrahydrofuranyl
138		NH ₂	H	neopentyl	2,5-pyrrolidinyl
139		NH ₂	H	neopentyl	3,4-dihydroxy-2,5-tetrahydrofuranyl
140		NH ₂	H	neopentyl	2,4-furanyl
141		NH ₂	H	neopentyl	4,2-furanyl
142		NH ₂	H	neopentyl	2,5-thienyl
143		NH ₂	H	neopentyl	2,4-thienyl
144		NH ₂	H	neopentyl	4,2-thienyl
145		NH ₂	H	neopentyl	2,5-pyrrolyl
146		NH ₂	H	neopentyl	2,5-imidazolyl
147		NH ₂	H	neopentyl	5,2-imidazolyl
148		NH ₂	H	neopentyl	2,5-oxazolyl
149		NH ₂	H	neopentyl	5,2-oxazolyl
150	12.1	NH ₂	H	neopentyl	3,4-dichloro-2,5-furanyl
151		NH ₂	H	neopentyl	3-chloro-2,5-furanyl
152		NH ₂	H	neopentyl	4-chloro-2,5-furanyl
153		NH ₂	H	neopentyl	3,4-fluoro-2,5-furanyl
154		NH ₂	H	neopentyl	3-fluoro-2,5-furanyl
155		NH ₂	H	neopentyl	4-fluoro-2,5-furanyl
156		NH ₂	H	neopentyl	CONHCH(CO ₂ H)
157		NH ₂	Me	neopentyl	2,5-furanyl
158		NH ₂	Et	neopentyl	2,5-furanyl
159		NH ₂	Pr	neopentyl	2,5-furanyl
160		NH ₂	vinyl	neopentyl	2,5-furanyl
161		NH ₂	acetylenyl	neopentyl	2,5-furanyl
162		NH ₂	allyl	neopentyl	2,5-furanyl
163		NH ₂	2-furanyl	neopentyl	2,5-furanyl
164		NH ₂	3-furanyl	neopentyl	2,5-furanyl
165		NH ₂	2-thienyl	neopentyl	2,5-furanyl
166		NH ₂	3-thienyl	neopentyl	2,5-furanyl
167		NH ₂	Ph	neopentyl	2,5-furanyl
168	22.1	NH ₂	NH ₂	neopentyl	2,5-furanyl
169		NH ₂	NHMe	neopentyl	2,5-furanyl
170		NH ₂	N(Me) ₂	neopentyl	2,5-furanyl

171		NH2	NHBn	neopentyl	2,5-furanyl
172		NH2	I	neopentyl	2,5-furanyl
173		NH2	Br	neopentyl	2,5-furanyl
174		NH2	Cl	neopentyl	2,5-furanyl
175		NH2	F	neopentyl	2,5-furanyl
176		NH2	OMe	neopentyl	2,5-furanyl
177		NH2	OEt	neopentyl	2,5-furanyl
178		NH2	OPr	neopentyl	2,5-furanyl
179		NH2	SO2Me	neopentyl	2,5-furanyl
180		NH2	SEt	neopentyl	2,5-furanyl
181		NH2	SPr	neopentyl	2,5-furanyl
182		NH2	SBu	neopentyl	2,5-furanyl
183		NH2	CN	neopentyl	2,5-furanyl
184		NH2	CONH2	neopentyl	2,5-furanyl
185		NH2	2-pyridyl	neopentyl	2,5-furanyl
186		NH2	3-pyridyl	neopentyl	2,5-furanyl
187		NH2	4-pyridyl	neopentyl	2,5-furanyl
188	5.4	NH2	H	1-(3-cyclohexyl)propyl	CH2OCH2
189	5.3	NH2	H	1-nonyl	CH2OCH2
190	5.2	NH2	H	2-cyclohexylethyl	CH2OCH2
191	5.1	NH2	H	2-phenethyl	CH2OCH2
192	10.2	NHMe	H	2-phenethyl	2,5-furanyl
193	10.1	N(Me)2	H	2-phenethyl	2,5-furanyl
194	9.1	Cl	H	2-phenethyl	2,5-furanyl
195	11.1	NH2	SMe	isobutyl	2,5-furanyl
196	11.2	NH2	SO2Me	isobutyl	2,5-furanyl
197	4.1	NH2	H	D-ribosyl	NHCH2CH2
198	4.2	NH2	H	5'-deoxy-D-ribosyl	NHCH2CH2
199		NH2	H	H	NHCH2CH2
200	3.1	NH2	H	benzyl	NHCH2CH2
201	3.2	NH2	H	2-phenethyl	NHCH2CH2
202	3.3	NH2	H	2-naphthylmethyl	NHCH2CH2
203	6.2	NH2	H	2-phenethyl	CH2CH2CH2
204	3.4	NH2	H	2-cyclohexylethyl	NHCH2CH2
205	6.1	NH2	H	2-cyclohexylethyl	CH2CH2CH2
206	8.1	NH2	H	2-cyclohexylethyl	SCH2
207	7.1	NH2	H	2-phenethyl	2,5-thienyl
208		NH2	Me	isobutyl	2,5-furanyl
209		NH2	Et	isobutyl	2,5-furanyl
210		NH2	SEt	isobutyl	2,5-furanyl
211		NH2	SPr	isobutyl	2,5-furanyl
212		NH2	2-furanyl	isobutyl	2,5-furanyl
213		NH2	2-thienyl	isobutyl	2,5-furanyl
214		NH2	Pr	isobutyl	2,5-furanyl

215		NH ₂	F	isobutyl	2,5-furanyl
216		NH ₂	Cl	isobutyl	2,5-furanyl
217		NH ₂	Br	isobutyl	2,5-furanyl
218		NH ₂	H	isobutyl	2,5-furanyl
219		NH ₂	Et	isobutyl	CONHCH ₂
220		NH ₂	SEt	isobutyl	CONHCH ₂
221		NH ₂	SPr	isobutyl	CONHCH ₂
222		NH ₂	2-furanyl	isobutyl	CONHCH ₂
223		NH ₂	2-thienyl	isobutyl	CONHCH ₂
224		NH ₂	Pr	isobutyl	CONHCH ₂
225		NH ₂	F	isobutyl	CONHCH ₂
226		NH ₂	Cl	isobutyl	CONHCH ₂
227		NH ₂	Br	isobutyl	CONHCH ₂
228		NH ₂	Me	isobutyl	CONHCH ₂
229		NH ₂	H	isobutyl	CONHCH ₂
230		NH ₂	Et	neopentyl	acetylene
231		NH ₂	SEt	neopentyl	acetylene
232		NH ₂	SPr	neopentyl	acetylene
233		NH ₂	2-furanyl	neopentyl	acetylene
234		NH ₂	2-thienyl	neopentyl	acetylene
235		NH ₂	Pr	neopentyl	acetylene
236		NH ₂	F	neopentyl	acetylene
237		NH ₂	Cl	neopentyl	acetylene
238		NH ₂	Br	neopentyl	acetylene
239		NH ₂	Me	neopentyl	acetylene
240		NH ₂	Et	neopentyl	NHCOCH ₂
241		NH ₂	SEt	neopentyl	NHCOCH ₂
242		NH ₂	SPr	neopentyl	NHCOCH ₂
243		NH ₂	2-furanyl	neopentyl	NHCOCH ₂
244		NH ₂	2-thienyl	neopentyl	NHCOCH ₂
245		NH ₂	Pr	neopentyl	NHCOCH ₂
246		NH ₂	F	neopentyl	NHCOCH ₂
247		NH ₂	Cl	neopentyl	NHCOCH ₂
248		NH ₂	Br	neopentyl	NHCOCH ₂
249		NH ₂	Me	neopentyl	NHCOCH ₂
250		NH ₂	Et	neopentyl	CH ₂ OCH ₂
251		NH ₂	SEt	neopentyl	CH ₂ OCH ₂
252		NH ₂	SPr	neopentyl	CH ₂ OCH ₂
253		NH ₂	2-furanyl	neopentyl	CH ₂ OCH ₂
254		NH ₂	2-thienyl	neopentyl	CH ₂ OCH ₂
255		NH ₂	Pr	neopentyl	CH ₂ OCH ₂
256		NH ₂	F	neopentyl	CH ₂ OCH ₂

257		NH ₂	Cl	n p ntyl	CH ₂ OCH ₂
258		NH ₂	Br	neopentyl	CH ₂ OCH ₂
259		NH ₂	Me	neopentyl	CH ₂ OCH ₂
260		NHBn	H	neopentyl	2,5-furanyl
261		NHPh	H	neopentyl	2,5-furanyl
262		NHBn	SMe	neopentyl	2,5-furanyl
263		NHPh	SMe	neopentyl	2,5-furanyl
264		NHPh-4-F	H	neopentyl	2,5-furanyl
265		NHPh-4-F	SMe	neopentyl	2,5-furanyl
266		NHNH ₂	F	neopentyl	2,5-furanyl
267		NH ₂	Me	cyclopropylmethyl	2,5-furanyl
268		NH ₂	SMe	cyclopropylmethyl	2,5-furanyl
269		NH ₂	F	cyclopropylmethyl	2,5-furanyl
270		NH ₂	Cl	cyclopropylmethyl	2,5-furanyl
271		NH ₂	Br	cyclopropylmethyl	2,5-furanyl
272		NH ₂	Et	cyclopropylmethyl	2,5-furanyl
273		NH ₂	CN	cyclopropylmethyl	2,5-furanyl
274		NH ₂	Me	cyclopropylmethyl	CONHCH ₂
275		NH ₂	SMe	cyclopropylmethyl	CONHCH ₂
276		NH ₂	F	cyclopropylmethyl	CONHCH ₂
277		NH ₂	Cl	cyclopropylmethyl	CONHCH ₂
278		NH ₂	Br	cyclopropylmethyl	CONHCH ₂
279		NH ₂	Et	cyclopropylmethyl	CONHCH ₂
280		NH ₂	CN	cyclopropylmethyl	CONHCH ₂
281		NH ₂	Me	cyclopropylmethyl	NHCOCH ₂
282		NH ₂	SMe	cyclopropylmethyl	NHCOCH ₂
283		NH ₂	F	cyclopropylmethyl	NHCOCH ₂
284		NH ₂	Cl	cyclopropylmethyl	NHCOCH ₂
285		NH ₂	Br	cyclopropylmethyl	NHCOCH ₂
286		NH ₂	Et	cyclopropylmethyl	NHCOCH ₂
287		NH ₂	CN	cyclopropylmethyl	NHCOCH ₂
288	2.18	NH ₂	H	3-(1-imidazolylpropyl)	2,5-furanyl
289	19.1	NH ₂	H	neopentyl	1,2-C ₆ H ₄ -O-
290	21.1	NH ₂	H	2-phenethyl	CONHCH ₂

More preferred are the following compounds from Table 1 including salts and prodrugs thereof:

- 5 1, 21, 22, 23, 29, 50, 57, 61, 62, 63, 66, 67, 72, 73, 89, 90, 107, 110, 111, 112, 113, 114, 115, 119, 123, 125, 126, 129, 130, 131, 132, 133, 134, 136, 137, 145, 148, 149, 151, 152, 153, 154, 155, 156, 158, 159, 163, 165, 173, 174, 175, 180, 181, 182, 183, 209, 210, 212, 215, 216, 217, 219, 220, 221, 230, 231, 234, 236, 237, 238, 240, 241, 246, 247, 248, 250, 251, 256, 257, 258, 266, 268, 269, and
10 272.

Most preferred are the following compounds and their salts and prodrugs:


- 5 N⁹-neopentyl-2-methylthio-8-phosphonomethylaminocarbonyladenine;
N⁹-neopentyl-2-methylthio-8-(2-(5-phosphono)furanyl)adenine;
N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine;
10 N⁹-isobutyl-2-methylthio-8-(2-(5-phosphono)furanyl)adenine;
N⁹-isobutyl-8-(2-(5-phosphono)furanyl)adenine;
N⁹-cyclopropyl-8-(2-(5-phosphono)furanyl)adenine;
15 N⁹-(2-cyclohexyl)ethyl-8-(2-(5-phosphono)furanyl)adenine;
N⁹-(1-(2,2-dimethyl-3-chloro)propyl)-8-(2-(5-phosphono)furanyl)adenine;
N⁹-(1-(3,3-dimethyl)butyl)-8-(2-(5-phosphono)furanyl)adenine;
20 N⁹-(1,5,5-trimethyl-3-cyclohexen-1-yl)methyl-8-(2-(5-phosphono)furanyl)adenine;
N⁹-neopentyl-8-(2-phosphonoacetylene-1-yl)adenine;
25 N⁹-neopentyl-8-(1-(3-phosphono-3-sulfuryl)propyl)adenine;
N⁹-neopentyl-8-(1-(3-phosphono-3-carboxyl)propyl)adenine;
30 N⁹-neopentyl-8-(1-(3,3-diphosphono)propyl)adenine;
N⁹-neopentyl-2-chloro-8-(2-(5-phosphono)furanyl)adenine;
2-Ethyl-N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine;
2-Methylthio-N⁹-isobutyl-8-(2-(5-phosphono)furanyl)adenine; and
2-Methylthio-N⁹-isobutyl-8-(phosphonomethoxymethyl)adenine.

35

Synthesis of Compounds of Formula 1

- Synthesis of compounds encompassed by the present invention typically includes some or all of the following general steps: (1) preparation of phosphonate prodrug; (2) deprotection of phosphonate ester; (3) modification
40 of C-8-substituted purine intermediates; (4) modification of purine at positions

other than C-8; (5) construction of the purine ring system; and (6) preparation of 4,5-diaminopyrimidine and other coupling partners.



5

(1) Preparation of phosphonate prodrugs

- Prodrug esters can be introduced at different stages of the synthesis. Because of their lability, prodrugs are often prepared from compounds of formula 1 where R¹ is H. Advantageously, these prodrug esters can be introduced at an early stage, provided that it can withstand the reaction conditions of the subsequent steps.
- Compounds of formula 5 where R¹ is H, can be alkylated with electrophiles (such as alkyl halides, alkyl sulfonates etc) under nucleophilic substitution reaction conditions to give phosphonate esters. For example prodrugs of formula 1, where R¹ is acyloxymethyl group can be synthesized through direct alkylation of the free phosphonic acid of formula 5, with the desired acyloxymethyl halide (e.g. Cl, Br, I; Elhaddadi, et al *Phosphorus Sulfur*, 1990, 54(1-4): 143; Hoffmann, *Synthesis*, 1988, 62) in presence of base e.g. *N, N'*-dicyclohexyl-4-morpholinecarboxamidine, Hunigs base etc. in polar aprotic solvents such as DMF (Starrett, et al, *J. Med. Chem.*, 1994, 1857). These carboxylates include but not limited to acetate, propylate, isobutyrate, pivalate, benzoate, and other carboxylates. Alternately, these acyloxymethylphosphonate esters can also be synthesized by treatment of the nitrophosphonic acid (A is NO₂ in formula 5; Dickson, et al, *J. Med. Chem.*, 1996, 39: 661; Iyer, et al, *Tetrahedron Lett.*, 1989, 30: 7141; Srivastva, et al, *Bioorg. Chem.*, 1984, 12: 118). This can be extended to many other types of

prodrugs, such as compounds of formula 1 where R¹ is 3-phthalidyl, 2-oxo-4,5-didehydro-1,3-dioxolanemethyl, and 2-oxotetrahydrofuran-5-yl groups, etc. (Biller and Magnin (US 5,157,027); Serafinowska et al. (*J. Med. Chem.* 38: 1372 (1995)); Starrett et al. (*J. Med. Chem.* 37: 1857 (1994)); Martin et al. *J. Pharm. Sci.* 76: 180 (1987); Alexander et al., *Collect. Czech. Chem. Commun.* 59: 1853 (1994)); and EPO 0632048A1). *N,N*-Dimethylformamide dialkyl acetals can also be used to alkylate phosphonic acids (Alexander, P., et al *Collect. Czech. Chem. Commun.*, 1994, 59, 1853).


Alternatively, these phosphonate prodrugs or phosphoramidates can also be synthesized, by reaction of the corresponding dichlorophosphonates and an alcohol or an amine (Alexander, et al, *Collect. Czech. Chem. Commun.*, 1994, 59: 1853). For example, the reaction of dichlorophosphonate with phenols and benzyl alcohols in the presence of base (such as pyridine, triethylamine, etc) yields compounds of formula 1 where R¹ is aryl (Khamnei, S., et al *J. Med. Chem.*, 1996, 39: 4109; Serafinowska, H.T., et al *J. Med. Chem.*, 1995, 38: 1372; De Lombaert, S., et al *J. Med. Chem.*, 1994, 37: 498) or benzyl (Mitchell, A.G., et al *J. Chem. Soc. Perkin Trans. 1*, 1992, 38: 2345). The disulfide-containing prodrugs, reported by Puech et al., *Antiviral Res.*, 1993, 22: 155, can also be prepared from dichlorophosphonate and 2-hydroxyethyl disulfide under the standard conditions.

Such reactive dichlorophosphonate intermediates, can be prepared from the corresponding phosphonic acids and the chlorinating agents e.g. thionyl chloride (Starrett, et al, *J. Med. Chem.*, 1994, 1857), oxalyl chloride (Stowell, et al, *Tetrahedron Lett.*, 1990, 31: 3261), and phosphorus pentachloride (Quast, et al, *Synthesis*, 1974, 490). Alternatively, these dichlorophosphonates can also be generated from disilyl phosphonate esters (Bhongle, et al, *Synth. Commun.*, 1987, 17: 1071) and dialkyl phosphonate esters (Still, et al, *Tetrahedron Lett.*, 1983, 24: 4405; Patois, et al, *Bull. Soc. Chim. Fr.*, 1993, 130: 485).

Furthermore, these prodrugs can be prepared from Mitsunobu reactions (Mitsunobu, *Synthesis*, 1981, 1; Campbell, *J. Org. Chem.*, 1992, 52: 6331), and other acid coupling reagents include, but not limited to, carbodiimides (Alexander, et al, *Collect. Czech. Chem. Commun.*, 1994, 59: 1853; Casara, et al, *Bioorg. Med. Chem. Lett.*, 1992, 2: 145; Ohashi, et al, *Tetrahedron Lett.*, 1988, 29: 1189), and benzotriazolyloxytris-(dimethylamino)phosphonium salts (Campagne, et al, *Tetrahedron Lett.*, 1993, 34: 6743). The prodrugs of formula

1 where R¹ is the cyclic carbonate or lactone or phthalidyl can also be synthesized by direct alkylation of free phosphonic acid with desired halides in the presence of base such as NaH or diisopropylethylamine (Biller and Magnin US 5,157,027; Serafinowska et al. *J. Med. Chem.* 38: 1372 (1995); Starrett et al. *J. Med. Chem.* 37: 1857 (1994); Martin et al. *J. Pharm. Sci.* 76: 180 (1987); Alexander et al., *Collect. Czech. Chem. Commun.* 59: 1853 (1994); and EPO 0632048A1).

R¹ can also be introduced at an early stage of synthesis, when feasible. For example, compounds of formula 1 where R¹ is phenyl can be prepared by phosphorylation of 2-furanylpurines via strong base treatment (e.g. LDA) followed by chlorodiphenylphosphonate, as shown in the following scheme. Alternatively, such compounds can be prepared by cyclization of 5-diphenylphosphono-2-furaldehyde with 4,5-diaminopyrimidines as described in section 5.



It is envisioned that compounds of formula 1 can be mixed phosphonate esters by combining the above described prodrugs (e.g. phenyl benzyl phosphonate esters, phenyl acyloxyalkyl phosphonate esters, etc.). For example, the chemically combined phenyl-benzyl prodrugs are reported by Meier et al. *Bioorg. Med. Chem. Lett.*, 1997, 7: 99.

The substituted cyclic propyl phosphonate esters of formula 5, can be synthesized by reaction of the corresponding dichlorophosphonate and the substituted 1,3-propanediol. The following are some of the methods to prepare the substituted 1,3-propanediols.


Synthesis of the 1,3-Propanediols Used in the Preparation of Certain Prodrugs

The discussion of this step includes various synthetic methods for the preparation of the following types of propane-1,3-diols: i) 1-substituted; ii) 2-substituted; and iii) 1,2- or 1,3-annulated. Different groups on the prodrug part of

th molecule *i.e.*, on the propanediol moiety can be introduced or modified either during the synthesis of the diols or after the synthesis of the prodrugs.


i) 1-Substituted 1,3-Propanediols

- 5 Propane-1,3-diols can be synthesized by several well known methods in the literature. Aryl Grignard additions to 1-hydroxypropan-3-al gives 1-aryl-substituted propane-1,3-diols (path a). This method will enable conversion of various substituted aryl halides to 1-arylsubstituted-1,3-propanediols (Coppi, et. al., *J. Org. Chem.*, 1988, 53, 911). Aryl halides can also be used to synthesize
- 10 1-substituted propanediols by Heck coupling of 1,3-diox-4-ene followed by reduction and hydrolysis (Sakamoto, et. al., *Tetrahedron Lett.*, 1992, 33, 6845). A variety of aromatic aldehydes can be converted to 1-substituted-1,3-propanediols by vinyl Grignard addition followed by hydroboration (path b). Substituted aromatic aldehydes are also utilized by lithium-t-butylacetate
- 15 addition followed by ester reduction (path e) (Turner., *J. Org. Chem.*, 1990, 55 4744). In another method, commercially available cinnamyl alcohols can be converted to epoxy alcohols under catalytic asymmetric epoxidation conditions. These epoxy alcohols are reduced by Red-Al to result in enantiomerically pure propane-1,3-diols (path c). Alternatively, enantiomerically pure 1,3-diols can be
- 20 obtained by chiral borane reduction of hydroxyethyl aryl ketone derivatives (Ramachandran, et. al., *Tetrahedron Lett.*, 1997, 38 761). Pyridyl, quinoline, and isoquinoline propan-3-ol derivatives can be oxygenated to 1-substituted propan-1,3-diols by N-oxide formation followed by rearrangement under acetic anhydride conditions (path d) (Yamamoto, et. al., *Tetrahedron*, 1981, 37, 1871).



ii) 2-Substituted 1,3-Propanediols:

- 5 Various 2-substituted propane-1,3-diols can be made from commercially available 2-(hydroxymethyl)-1,3-propanediol. Triethyl methanetricarboxylate can be converted to the triol by complete reduction (path a) or diol-monocarboxylic acid derivatives can be obtained by partial hydrolysis and diester reduction (Larock, *Comprehensive Organic Transformations*, VCH, New York, 1989). Nitrotriol is also known to give the triol by reductive elimination (path b) (Latour, et. al., *Synthesis*, 1987, 8, 742). The triol can be derivatized as a mono acetate or carbonate by treatment with alkanoyl chloride, or alkylchloroformate, respectively (path d) (Greene and Wuts, *Protective Groups in Organic Synthesis*, John Wiley, New York, 1990). Aryl substitution can be made by oxidation to the aldehyde followed by aryl Grignard additions (path c) and the aldehyde can also be converted to substituted amines by reductive amination reactions (path e).
- 10
- 15



iii) Annulated 1,3-Propanediols:

Prodrugs of formula 1 where V - Z or V - W are fused by three carbons are made from cyclohexanediol derivatives. Commercially available *cis*, *cis*-1,3,5-cyclohexanetriol can be used for prodrug formation. This cyclohexanetriol can also be modified as described in the case of 2-substituted propane-1,3-diols to give various analogues. These modifications can either be made before or after formation of prodrugs. Various 1,3-cyclohexanediols can be made by

Diels-Alder methodology using pyrone as the diene (Posner, et. al., *Tetrahedron Lett.*, **1991**, *32*, 5295). Cyclohexyl diol derivatives are also made by nitrile oxide olefin-additions (Curran, et. al., *J. Am. Chem. Soc.*, **1985**, *107*, 6023). Alternatively, cyclohexyl precursors can be made from quinic acid (Rao, et. al., *Tetrahedron Lett.*, **1991**, *32*, 547.)

(2) Deprotection of Phosphonate Esters

Compounds of formula 1 where R¹ is H may be prepared from phosphonate esters using known phosphate and phosphonate ester cleavage conditions. For example, alkyl phosphonate esters are generally cleaved by reaction with silyl halides followed by hydrolysis of the intermediate silyl phosphonate esters. Depending on the stability of the products, these reactions are usually accomplished in the presence of acid scavengers such as

1,1,1,3,3,3-hexamethyldisilazane, 2,6-lutidine, etc. Various silyl halides can be used for this transformation, such as chlorotrimethylsilane (Rabinowitz *J. Org. Chem.*, **1963**, *28*: 2975), bromotrimethylsilane (McKenna et al. *Tetrahedron Lett.*, **1977**, 155), iodotrimethylsilane (Blackburn et al. *J. Chem. Soc., Chem. Commun.*, **1978**, 870). Phosphonate esters can also be cleaved under strong acid conditions, such as hydrogen halides in acetic acid or water, and metal halides (Moffatt et al. *U.S. Patent 3,524,846*, **1970**). Phosphonate esters can also be converted to dichlorophosphonates with halogenating agents (e.g. PCl_5 , SOCl_2 , BBr_3 , etc. Pelchowicz et al. *J. Chem. Soc.*, **1961**, 238) and subsequently hydrolyzed to give phosphonic acids. Reductive reactions are useful in cleaving aryl and benzyl phosphonate esters. For example, aryl and benzyl phosphonate esters can be cleaved under hydrogenolysis conditions (Lejczak et al. *Synthesis*, **1982**, 412; Elliott et al. *J. Med. Chem.*, **1985**, *28*: 1208.) or dissolving metal reduction conditions (Shafer et al. *J. Am. Chem. Soc.*, **1977**, *99*: 5118). (Elliott et al. *J. Med. Chem.*, **1985**, *28*: 1208). Electrochemical (Shono et al. *J. Org. Chem.*, **1979**, *44*: 4508) and pyrolysis (Gupta et al. *Synth. Commun.*, **1980**, *10*: 299) conditions have also been used to cleave various phosphonate esters.

(3) Modification of C-8-substituted Purine Intermediates

8-Substituted purines are useful intermediates in the preparation of compounds of formula 1. 8-Halopurines, which are particularly useful intermediates, are readily prepared using chemistry well described in the literature. For example, N^9 -alkyladenines are halogenated at the C-8 position using known halogenating agents (e.g. Br_2 , NBS). 8-Alkylpurine can be prepared through direct lithiation of purine followed by trapping with electrophiles (e.g. alkyl halides, Barton et al. *Tetrahedron Lett.*, **1979**, 5877).

Functionalization of 8-halopurines can be accomplished under substitution reaction conditions with nucleophiles such as amines, alcohols, azides, sulfides, and alkylthiols. It is advantageous to have the phosphonate moiety as part of the nucleophiles. For example, substitution of 8-bromopurine with aminoalkylphosphonates gives compounds of formula 1 where X is alkylamino.

8-Halopurines can also be transformed into other 8-substituted purines using palladium catalyzed reactions (Heck *Palladium Reagents in Organic Synthesis*; Academic Press: San Diego, 1985). For example, palladium catalyzed carbonylation reactions of 8-bromopurine in the presence of alcohol
5 gives 8-alkoxycarbonyl purines. Using known chemistry, the 8-carboxylate group can be converted into other functional groups, such as hydroxymethyl, halomethyl, formyl, carboxylic acid, carbamoyl, and thiocarbonyl groups. These functional groups are useful intermediates for the synthesis of compounds of formula 1. For example, 8-alkyl and 8-aryl purines can be prepared from 8-
10 halopurines via palladium catalyzed coupling reactions with organotin (Moriarty et al. *Tetrahedron Lett.*, 1990, 41: 5877), organoborane (Yatagai, *Bull. Chem. Soc. Jpn.*, 1980, 53: 1670), and other reagents known to couple with aryl halides. When the coupling reagents contain the dialkylphosphonate group, the reaction is useful for preparation of compounds of formula 5 where X is alkyl,
15 alkenyl, alkynyl, and aryl. For example, 8-bromopurine can be coupled with diethyl 1-tributylstannyl-3-allylphosphonate to give compounds of formula 5 where X is $-\text{CH}=\text{CHCH}_2-$. Subsequent hydrogenation reaction gives compounds of formula 5 where X is $-\text{CH}_2\text{CH}_2\text{CH}_2-$.

The phosphonate group can also be introduced by further modification of
20 the 8-substituents. Substitutions of 8-haloalkyl or 8-sulfonylalkyl purine with nucleophiles containing the phosphonate group are useful for the preparation of compounds of formula 5 where X is alkylaminoalkyl, alkoxyalkyl, and alkylthioalkyl. For example, compounds of formula 5 where X is $-\text{CH}_2\text{OCH}_2-$ can be prepared from 8-bromomethylpurine using hydroxymethylphosphonate
25 esters and a suitable base. It is possible to reverse the nature of the nucleophiles and electrophiles for the substitution reactions, i.e. haloalkyl- and/or sulfonylalkylphosphonate esters can be substituted with purines containing a nucleophile at the C-8 position (such as 8-hydroxyalkyl, 8-thioalkyl, and 8-aminoalkyl purines). For example, diethyl phosphonomethyltriflate can be
30 substituted by alcohols such as 8-hydroxymethylpurine to give compounds of formula 5 where X is $-\text{CH}_2\text{OCH}_2-$ (Phillion et al. *Tetrahedron Lett.* 1986, 27: 1477). Known amide formation reactions are useful for the synthesis of compounds of formula 5 where X is alkylaminocarbonyl, alkoxy carbonyl, alkoxythiocarbonyl, and alkylthiocarbonyl. For example, coupling of 8-
35 purinecarboxylic acids with aminoalkylphosphonate esters gives compounds of formula 5 where X is alkylaminocarbonyl. For compounds of formula 5 where X

is alkyl, the phosphonate group can also be introduced using other common phosphonate formation methods, such as Michaelis-Arbuzov reaction (Bhattacharya et al. *Chem. Rev.*, 1981, 81: 415), Michaelis-Becker reaction (Blackburn et al. *J. Organomet. Chem.*, 1988, 348: 55), addition reactions of phosphorus to electrophiles (such as aldehydes, ketones, acyl halides, imines and other carbonyl derivatives).

8-Azidopurines are useful for the preparation for compounds of formula 5 where X is alkylamino and alkylcarbonylamino groups. For example, carboxylic acids (e.g. $(RO)_2P(O)$ -alkyl-CO₂H) can be directly coupled to 8-azidopurines to give 8-alkylcarbonylaminopurines (Urpi et al. *Tetrahedron Lett.*, 1986, 27: 4623). Alternatively, 8-azidopurines can also be converted to 8-aminopurines under reductive conditions, and subsequently converted to 8-alkylaminocarbonyl- and 8-alkylaminopurines using known chemistry.

(4) Modification of purines at positions other than C-8

Compounds of formula 5 can be further modified to give intermediates useful for the synthesis of compounds of formula 1. For example, substitution reactions of 6-chloropurine by ammonia or alkylamines are useful for the preparations of compounds of formula 5 where A is amino and alkylamino groups.

E groups can be introduced by modifying existing 2-substituents of purine. For example, 2-halopurines, readily accessible from 2-aminopurines via chemistry well described in the literature, can be converted to other 2-substituted purines by, for example, nucleophilic substitution reactions; transition metal catalyzed reactions, etc. (*J. Med. Chem.*, 1993, 36: 2938; *Heterocycles*, 1990, 30: 435).

E groups can also be introduced via metalation (e.g. lithiation, *J. Org. Chem.*, 1997, 62(20), 6833) of the C-2-position and followed by addition of electrophiles which can be the desired E group or a substituent (e.g. tributylstannyl group) which can be converted to the desired E group using conventional chemistry.

It is envisioned that N⁹-substituted purines can be readily prepared from compounds of formula 5 where Y is H using, for example, standard alkylation reactions (with alkyl halide, or sulfonate), or Mitsunobu reactions. Further elaborations of substituents on Y are also possible.


More importantly, combinatorial methods have been developed for synthesis of 2- and N-9-substituted purines on solid-phase which conceivably can be applied for the synthesis of purine FBPase inhibitors (Schultz, et al, *Tetrahedron Lett.*, 1997, 38(7), 1161; *J. Am. Chem. Soc.*, 1996, 118, 7430).

5

(5) Construction of the Purine Ring System

The purine ring system of compounds of formula 1 can be constructed using 4,5-diaminopyrimidines and carboxylates or their derivatives (such as aldehydes, amides, nitriles, ortho esters, imidates, etc.) (Townsend *Chemistry of Nucleosides and Nucleotides, Vol 1* ; Plenum Press, New York and London, page 156-158). For example, alkyl and aryl aldehydes can be cyclized with 4,5-

10



Intramolecular cyclization reactions of pyrimidine derivatives can also be used to construct the purine ring system. For example, 5-acylamino-4-alkylaminopyrimidines are treated with phosphorus oxychloride and cyclized under basic conditions to give purine derivatives. This transformation can also be achieved using other reagents (e.g. SiCl₄-Et₃N, Desaubry et al. *Tetrahedron Lett.*, 1995, 36: 4249). Imidazole derivatives are also useful for the construction of purine ring system via cyclization reactions to form the pyrimidine ring (Townsend *Chemistry of Nucleosides and Nucleotides, Vol 1* ; Plenum Press, New York and London, page 148-156).

(6) Preparation of Diaminopyrimidine and Other Coupling Partners

Compounds of formula 4 are useful for the construction of purine ring systems, and such compounds can be readily synthesized using known chemistry. For example, the Y group can be introduced using a nucleophilic substitution reaction involving an amine and 4-halopyrimidines (*Tetrahedron*, 1984, 40: 1433). Alternatively, palladium catalyzed reactions (Wolfe et al. *J. Am. Chem. Soc.*, 1996, 118: 7215) can also be used. Reductive amination reactions (*Synthesis*, 1975, 135) and alkylation with electrophiles (such as

30

halides, sulfonates) are useful for the preparation of compounds of formula 4 from 4-aminopyrimidines. The 5-amino group can be introduced using amine formation reactions such as nitration followed by reduction (Dhainant et al. *J. Med. Chem.*, 1996, 39: 4099), arylazo compound formation followed by
5 reduction (Lopez et al. *Nucleosides & Nucleotides*, 1996, 15: 1335), azide formation followed by reduction, or by rearrangement of carboxylic acid derivatives (e.g. Schmidt, Curtius, and Beckmann reactions).

Coupling of aromatic or aliphatic aldehydes, and carboxylic acid derivatives with attached phosphonate esters are particularly suited for the
10 preparation of compounds of formula 1 as described in section 5. Such phosphonate esters are prepared by lithiation of the aromatic ring using methods well described in literature (Gschwend *Org. React.* 1979, 26: 1) followed by addition of phosphorylating agents (e.g. ClPO_3R_2). Phosphonate esters can also be introduced by Arbuzov-Michaelis reaction (Brill *Chem Rev.*,
15 1984, 84: 577) and transition metal catalyzed reaction with alkyl halides and aryl halides or triflates (Balthazar et al. *J. Org. Chem.*, 1980, 45: 5425; Petrakis et al. *J. Am. Chem. Soc.*, 1987, 109: 2831; Lu et al. *Synthesis*, 1987, 726). Alternatively, aryl phosphonate esters can be prepared from aryl phosphates under anionic rearrangement conditions (Melvin *Tetrahedron Lett.*, 1981, 22:
20 3375; Casteel et al. *Synthesis*, 1991, 691). Aryl phosphate esters can also be used to prepare compounds of Formula 1 where X is an oxyaryl group. N-Alkoxy aryl salts with alkali metal derivatives of dialkyl phosphonate can be used to synthesize heteroaryl-2-phosphonate esters (Redmore *J. Org. Chem.*, 1970, 35: 4114).

25 A second lithiation step can be used to incorporate the aldehyde functionality, although other methods known to generate aromatic aldehydes can be envisioned as well (e.g. Vilsmeier-Hack reaction, Reimar-Teimann reaction etc.). In the second lithiation step, the lithiated aromatic ring is treated with reagents that either directly generate an aldehyde (e.g. DMF, HCO_2R , etc.)
30 or with reagents that lead to a group that subsequently is transformed into an aldehyde group using known chemistry (e.g. alcohol, ester, cyano, alkene, etc.). It is also envisioned that the sequence of these reactions can be reversed, i.e. the aldehyde moiety can be incorporated first followed by the phosphorylation reaction. The order of the reaction will depend on the reaction conditions and
35 the protecting groups. Prior to the phosphorylation, it is also envisioned that it may be advantageous to protect the aldehydes using a number of well-known

steps (hemiacetal, hemiaminal, etc.). The aldehyde is then unmasked after phosphorylation. (*Protective groups in Organic Synthesis*, Greene, T. W., 1991, Wiley, New York).

5 Formulations

Compounds of the invention are administered orally in a total daily dose of about 0.1 mg/kg/dose to about 100 mg/kg/dose, preferably from about 0.3 mg/kg/dose to about 30 mg/kg/dose. The most preferred dose range is from 0.5 to 10 mg/kg (approximately 1 to 20 nmoles/kg/dose). The use of time-release preparations to control the rate of release of the active ingredient may be preferred. The dose may be administered in as many divided doses as is convenient. When other methods are used (e.g. intravenous administration), compounds are administered to the affected tissue at a rate from 0.3 to 300 nmol/kg/min, preferably from 3 to 100 nmoles/kg/min. Such rates are easily maintained when these compounds are intravenously administered as discussed below.

For the purposes of this invention, the compounds may be administered by a variety of means including orally, parenterally, by inhalation spray, topically, or rectally in formulations containing pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used here includes subcutaneous, intravenous, intramuscular, and intraarterial injections with a variety of infusion techniques. Intraarterial and intravenous injection as used herein includes administration through catheters. Oral administration is generally preferred.

Pharmaceutical compositions containing the active ingredient may be in any form suitable for the intended method of administration. When used for oral use for example, tablets, troches, lozenges, aqueous or oil suspensions, dispersible powders or granules, emulsions, hard or soft capsules, syrups or elixirs may be prepared. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents including sweetening agents, flavoring agents, coloring agents and preserving agents, in order to provide a palatable preparation. Tablets containing the active ingredient in admixture with non-toxic pharmaceutically acceptable excipient which are suitable for manufacture of tablets are acceptable. These excipients may be, for example, inert diluents, such as calcium or sodium carbonate,

lactose, calcium or sodium phosphate; granulating and disintegrating agents, such as maize starch, or alginic acid; binding agents, such as starch, gelatin or acacia; and lubricating agents, such as magnesium stearate, stearic acid or talc. Tablets may be uncoated or may be coated by known techniques including
5 microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

Formulations for oral use may be also presented as hard gelatin
10 capsules where the active ingredient is mixed with an inert solid diluent, for example calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, such as peanut oil, liquid paraffin or olive oil.

Aqueous suspensions of the invention contain the active materials in
15 admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients include a suspending agent, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia, and dispersing or wetting agents such as a naturally occurring phosphatide (e.g.,
20 lecithin), a condensation product of an alkylene oxide with a fatty acid (e.g., polyoxyethylene stearate), a condensation product of ethylene oxide with a long chain aliphatic alcohol (e.g., heptadecaethyleneoxycetanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan monooleate). The aqueous
25 suspension may also contain one or more preservatives such as ethyl or n-propyl p-hydroxy-benzoate, one or more coloring agents, one or more flavoring agents and one or more sweetening agents, such as sucrose or saccharin.

Oil suspensions may be formulated by suspending the active ingredient in a vegetable oil, such as arachis oil, olive oil, sesame oil or coconut oil, or in a
30 mineral oil such as liquid paraffin. The oral suspensions may contain a thickening agent, such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

Dispersible powders and granules of the invention suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent, and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those disclosed above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, a mineral oil, such as liquid paraffin, or a mixture of these. Suitable emulsifying agents include naturally-occurring gums, such as gum acacia and gum tragacanth, naturally occurring phosphatides, such as soybean lecithin, esters or partial esters derived from fatty acids and hexitol anhydrides, such as sorbitan monooleate, and condensation products of these partial esters with ethylene oxide, such as polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, such as glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, a flavoring or a coloring agent.

The pharmaceutical compositions of the invention may be in the form of a sterile injectable preparation, such as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as a solution in 1,3-butane-diol or prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile fixed oils may conventionally be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables.

The amount of active ingredient that may be combined with the carrier material to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a time-release formulation intended for oral administration to humans may

contain 20 to 2000 μmol (approximately 10 to 1000 mg) of active material compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95% of the total compositions. It is preferred that the pharmaceutical composition be prepared which provides easily measurable amounts for administration. For example, an aqueous solution intended for intravenous infusion should contain from about 0.05 to about 50 μmol (approximately 0.025 to 25 mg) of the active ingredient per milliliter of solution in order that infusion of a suitable volume at a rate of about 30 mL/hr can occur.

As noted above, formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be administered as a bolus, electuary or paste.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free flowing form such as a powder or granules, optionally mixed with a binder (e.g., povidone, gelatin, hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (e.g., sodium starch glycolate, cross-linked povidone, cross-linked sodium carboxymethyl cellulose) surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropyl methylcellulose in varying proportions to provide the desired release profile. Tablets may optionally be provided with an enteric coating, to provide release in parts of the gut other than the stomach. This is particularly advantageous with the compounds of formula 1 when such compounds are susceptible to acid hydrolysis.

Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredient in a flavored base, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert

base such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

Formulations for rectal administration may be presented as a suppository with a suitable base comprising for example cocoa butter or a salicylate.

5 Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

Formulations suitable for parenteral administration include aqueous and
10 non-aqueous isotonic sterile injection solutions which may contain antioxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose sealed
15 containers, for example, ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

20 Preferred unit dosage formulations are those containing a daily dose or unit, daily sub-dose, or an appropriate fraction thereof, of a fructose 1,6-bisphosphatase inhibitor compound.

It will be understood, however, that the specific dose level for any particular patient will depend on a variety of factors including the activity of the
25 specific compound employed; the age, body weight, general health, sex and diet of the individual being treated; the time and route of administration; the rate of excretion; other drugs which have previously been administered; and the severity of the particular disease undergoing therapy, as is well understood by those skilled in the art.

30

Utility

FBPase inhibitors at the AMP site may be used to treat diabetes mellitus, lower blood glucose levels, and inhibit gluconeogenesis.

35 FBPase inhibitors at the AMP site may also be used to treat excess glycogen storage diseases. Excessive hepatic glycogen stores are found in patients with some glycogen storage diseases. Since the indirect pathway

contributes significantly to glycogen synthesis (Shulman, G.I. *Phys. Rev.* **1992**, *72*, 1019-1035), inhibition of the indirect pathway (gluconeogenesis flux) is expected to decrease glycogen overproduction.

FBPase inhibitors at the AMP site may also be used to treat or prevent
5 diseases associated with increased insulin levels.

Increased insulin levels are associated with an increased risk of cardiovascular complications and atherosclerosis (Folsom, et al., *Stroke*, **1994**, *25*, 66-73; Howard, G. et al., *Circulation* **1996**, *93*, 1809-1817). FBPase inhibitors are expected to decrease postprandial glucose levels by enhancing
10 hepatic glucose uptake. This effect is postulated to occur in individuals that are non-diabetic (or pre-diabetic, i.e. without elevated HGO or fasting blood glucose levels). Increased hepatic glucose uptake will decrease insulin secretion and thereby decrease the risk of diseases or complications that arise from elevated insulin levels.

15 The compounds of this invention and their preparation can be understood further by the examples which illustrate some of the processes by which these compounds are prepared. These examples should not, however, be construed as specifically limiting the invention and variations of the invention, now known or later developed, are considered to fall within the scope
20 of the present invention as hereinafter claimed.

EXAMPLES

Example 1

25 **Preparation of 5-diethylphosphono-2-furaldehyde (1).**

Step A. A solution of 2-furaldehyde diethyl acetal (1 mmol) in THF was treated with nBuLi (1 mmol) at -78 °C. After 1 h, diethyl chlorophosphate (1.2 mmol) was added and the reaction was stirred for 40 min. Extraction and
30 evaporation gave a brown oil.

Step B. The resulting brown oil was treated with 80 % acetic acid at 90 °C for 4 h. Extraction and chromatography gave compound 1 as a clear yellow oil.

Alternatively this aldehyde can be prepared from furan as described
35 below.

Step C. A solution of furan (1 mmol) in diethyl ether was treated with TMEDA (N,N,N',N'-tetramethylethylenediamine) (1 mmol) and nBuLi (2 mmol) at -78 °C. The solution was stirred for 0.5 h. at -78 °C and diethyl chlorophosphate was added and stirred for another 1 h. Extraction and distillation produced
5 diethyl 2-furanphosphonate as a clear oil.

Step D. A solution of diethyl 2-furanphosphonate (1 mmol) in THF (tetrahydrofuran) was treated with LDA (1.12 mmol, lithium N,N-diisopropylamide) at -78 °C for 20 min. Methyl formate (1.5 mmol) was added and the reaction was stirred for 1 h. Extraction and chromatography gave
10 compound **1** as a clear yellow oil.

Preferably this aldehyde can be prepared from 2-furaldehyde as described below.

Step E. A solution of 2-furaldehyde (1 mmol) and N,N'-dimethylethylene diamine (1 mmol) in toluene was refluxed with a Dean-Stark trap to collect the
15 resulting water. After 2 h the solvent was removed in vacuo and the residue was distilled to give furan-2-(N,N'-dimethylimidazolidine) as a clear colorless oil, bp 59-61 °C (3 mm Hg).

Step F. A solution of furan-2-(N,N'-dimethylimidazolidine) (1 mmol) and TMEDA (1 mmol) in THF was treated with nBuLi (1.3 mmol) at -40 to -48 °C.
20 The reaction was stirred at 0 °C for 1.5 h and then cooled to -55 °C and treated with a solution of diethylchlorophosphate (1.1 mmol) in THF. After stirring at 25 °C for 12 h the reaction mixture was evaporated and subjected to extraction to give 5-diethylphosphono-furan-2-(N,N'-dimethylimidazolidine) as a brown oil.

Step G. A solution of 5-diethylphosphonofuran-2-(N,N'-dimethylimidazolidine) (1 mmol) in water was treated with concentrated sulfuric acid
25 until pH = 1. Extraction and chromatography gave compound **1** as a clear yellow oil.

Example 2.

30 Preparation of N⁹-substituted-8-(2-(5-phosphono)furanyl)adenines.

The preparation of N⁹-(2-phenethyl)-8-(2-(5-phosphono)furanyl)adenine is given as an example:

Step A. A solution of 5-amino-4,6-dichloropyrimidine (1 mmol) in nBuOH was treated with Et₃N (1.2 mmol) and phenethylamine (1.05 mmol) at 80 °C.
35 After 12 h, the cooled reaction mixture was evaporated under vacuum and the residue was chromatographed to give 6-chloro-5-amino-4-(phenethylamino)-

pyrimidine as a yellow solid. mp 156 - 157 °C; TLC: R_f = 0.41, 50 % EtOAc-hexane.

Step B. The 6-chloro-5-amino-4-(2-phenethylamino)pyrimidine (1 mmol) in DMSO was treated with 2-furaldehyde (1.5 mmol) and FeCl_3 -silica (2.0 mmol) at 80 °C. After 12 h, the cooled reaction mixture was filtered and the filtrate was evaporated under vacuum. Chromatography afforded 6-chloro- N^9 -(2-phenethyl)-8-(2-furanyl)purine as a yellow solid. TLC: R_f = 0.62, 50 % EtOAc-hexane. Anal. calcd. for $\text{C}_{17}\text{H}_{13}\text{N}_4\text{OCl}$: C: 62.87; H: 4.03; N: 17.25. Found: C: 62.66; H: 3.96; N: 17.07.

Step C. The 6-chloro- N^9 -(2-phenethyl)-8-(2-furanyl)purine (1 mmol) in THF was treated with LDA (1.5 mmol) at -78 °C. After 1 h, diethyl chlorophosphate (5 mmol) was added and the reaction was stirred at -78 °C for 2 h and then quenched with saturated NH_4Cl . Extraction and chromatography gave 6-chloro- N^9 -(2-phenethyl)-8-(2-(5-diethylphosphono)-furanyl)purine as a yellow solid. TLC: R_f = 0.34, 100 % EtOAc.

Alternatively this type of compound can be prepared as follows:

Step D. A solution of 6-chloro-5-amino-4-(2-phenethylamino)pyrimidine (1 mmol) in DMSO was treated with 5-diethylphosphono-2-furaldehyde (1, 1.5 mmol), and FeCl_3 -silica (2.0 mmol) at 80 °C. After 12 h., the cooled reaction mixture was filtered and the filtrate was evaporated under vacuum. Chromatography afforded 6-chloro- N^9 -(2-phenethyl)-8-(2-(5-diethylphosphono)furanyl)purine as a yellow solid. TLC: R_f = 0.34, 100 % EtOAc.

Step E. 6-Chloro- N^9 -(2-phenethyl)-8-(2-(5-diethylphosphono)furanyl)purine (1 mmol) in THF-DMSO was treated with liquid ammonia (2 mL) in a steel bomb. After 12 h, the reaction was evaporated under vacuum and the residue was purified through chromatography to give N^9 -(2-phenethyl)-8-(2-(5-diethylphosphono)furanyl)adenine as a yellow solid. TLC: R_f = 0.12, 5 % MeOH - CH_2Cl_2 .

Step F. A solution of N^9 -(2-phenethyl)-8-(2-(5-diethylphosphono)furanyl)adenine (1 mmol) in acetonitrile was treated with bromotrimethylsilane (10 mmol). After 12 h, the reaction was evaporated under vacuum and the residue was treated with a mixture of water and acetonitrile. The solid was collected through filtration to give N^9 -(2-phenethyl)-8-(2-(5-phosphono)furanyl)adenine (2.1). mp 242-244° C; Anal. calcd. for $\text{C}_{17}\text{H}_{16}\text{N}_5\text{O}_4\text{P} + 1.37\text{H}_2\text{O}$: C: 50.16; H: 4.64; N: 17.21. Found: C: 48.95; H: 4.59; N: 16.80.

The following compounds were prepared according to above procedures:

- 2.2: N⁹-(2-cyclohexylethyl)-8-(2-(5-phosphono)furanyl)adenine. mp 194-195 °C; Anal. calcd. for C₁₇H₂₂N₅O₄P + 1 H₂O: C: 49.90; H: 5.90; N: 17.10. Found: C: 50.20; H: 5.70; N: 17.10.
- 2.3: N⁹-(2-naphthylmethyl)-8-(2-(5-phosphono)furanyl)adenine. mp 255-256 °C; Anal. calcd. for C₂₀H₁₆N₅O₄P + 1 H₂O: C: 54.70; H: 4.10; N: 15.90. Found: C: 54.30; H: 4.20; N: 15.90.
- 2.4: N⁹-(1-(2,2-diphenyl)ethyl)-8-(2-(5-phosphono)furanyl)adenine. mp 220-221 °C; Anal. calcd. for C₂₃H₂₀N₅O₄P + 0.25 H₂O: C: 59.29; H: 4.43; N: 15.03. Found: C: 59.35; H: 4.25; N: 14.83.
- 2.5: N⁹-ethyl-8-(2-(5-phosphono)furanyl)adenine. mp > 230 °C; Anal. calcd. for C₁₁H₁₂N₅O₄P + 1 H₂O: C: 40.38; H: 4.31; N: 21.40. Found: C: 40.45; H: 4.18; N: 21.44.
- 2.6: N⁹-isobutyl-8-(2-(5-phosphono)furanyl)adenine. mp > 230 °C; Anal. calcd. for C₁₃H₁₆N₅O₄P: C: 46.30; H: 4.78; N: 20.76. Found: C: 46.00; H: 4.61; N: 20.49.
- 2.7: N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine. mp > 230 °C; Anal. calcd. for C₁₄H₁₈N₅O₄P: C: 47.87; H: 5.16; N: 19.94. Found: C: 47.59; H: 4.92; N: 19.53.
- 2.8: N⁹-adamantanemethyl-8-(2-(5-phosphono)furanyl)adenine. mp > 250 °C; Anal. calcd. for C₂₀H₂₄N₅O₄P + 0.5 H₂O + 0.25 MeOH: C: 54.48; H: 5.87; N: 15.69. Found: C: 54.62; H: 5.52; N: 15.36.
- 2.9: N⁹-cyclopropyl-8-(2-(5-phosphono)furanyl)adenine. mp > 250 °C; MS (M + H) calcd for C₁₂H₁₃N₅O₄P: 322, found: 322.
- 2.10: N⁹-cyclopentyl-8-(2-(5-phosphono)furanyl)adenine. mp 220 °C(decomp); Anal. calcd. for C₁₄H₁₆N₅O₄P + 1 H₂O: C: 45.78; H: 4.94; N: 19.07. Found: C: 45.40; H: 4.79; N: 18.73.
- 2.11: N⁹-((2-ethoxy)phenyl)methyl-8-(2-(5-phosphono)furanyl)-adenine. mp > 230 °C; Anal. calcd. for C₁₈H₁₈N₅O₅P + 2 H₂O: C: 47.90; H: 4.91; N: 15.52. Found: C: 48.03; H: 4.53; N: 15.25.
- 2.12: N⁹-(1-(3-N,N-dimethylamino-2,2-dimethyl)propyl)-8-(2-(5-phosphono)furanyl)adenine. mp > 230 °C; Anal. calcd. for C₁₆H₂₃N₆O₄P + 3 H₂O + 0.5 HOAc + 0.75 Na: C: 41.19; H: 6.30; N: 16.95. Found: C: 41.35; H: 6.04; N: 16.57.

- 2.13:** N⁹-(1-(3-hydroxyl-2,2-dimethyl)propyl)-8-(2-(5-phosphono)-furanyl)adenine. mp > 230 °C; Anal. calcd. for C₁₄H₁₈N₅O₅P + 0.25 H₂O: C: 45.23; H: 5.02; N: 18.83. Found: C: 45.40; H: 5.02; N: 18.44.
- 2.14:** N⁹-(1-(3-chloro-2,2-dimethyl)propyl)-8-(2-(5-phosphono)-furanyl)adenine.
5 mp > 230 °C; Anal. calcd. for C₁₄H₁₇N₅O₄PCl + 0.125 CHCl₃ + 0.06 AcOEt: C: 42.50; H: 4.37; N: 17.25. Found: C: 42.62; H: 3.99; N: 16.87.
- 2.15:** N⁹-(1-(3,3-dimethyl)butyl)-8-(2-(5-phosphono)furanyl)-adenine. mp 230 °C; Anal. calcd. for C₁₅H₂₀N₅O₄P + 1.25H₂O + 0.13 AcOEt: C: 46.68; H: 5.94; N: 17.56. Found: C: 46.67; H: 5.78; N: 17.35.
- 10 **2.16:** N⁹-(1,5,5-trimethyl-3-cyclohexenyl)methyl-8-(2-(5-phosphono)furanyl)-adenine. mp > 230 °C; Anal. calcd. for C₁₉H₂₄N₅O₄P + 0.5 H₂O + 0.13 AcOEt: C: 53.54; H: 5.99; N: 16.01. Found: C: 53.67; H: 5.69; N: 15.75.
- 2.17:** N⁹-(1-(1,2,2-trimethyl)propyl)-8-(2-(5-phosphono)-furanyl)adenine. mp > 250 °C; Anal. calcd. for C₁₅H₂₀N₅O₄P + 0.67 H₂O + 0.13AcO Et: C: 47.74; H: 5.70;
15 N: 18.56. Found: C: 47.99; H: 5.39; N: 18.49.
- 2.18:** 6-Amino-9-(3-(1-imidazolyl)propyl)-8-(2-(5-phosphono)furanyl)purine. mp 182-186°C; Mass calcd. for C₁₅H₁₆N₇O₄P: 389. Found: M + H⁺ = 390.

Examples 3.

- 20 Preparation of N⁹-substituted-8-(2-phosphonoethylamino)adenines.
- Step A. Adenine (1 mmol) in DMF was treated with sodium hydride (1.2 mmol) followed by benzyl bromide (1.2 mmol) at room temperature under nitrogen. The resulting mixture was warmed at 100 °C for 2 h. The cooled reaction mixture was evaporated to dryness. Extraction and chromatography
25 afforded N⁹-benzyladenine.
- Step B. A solution of N⁹-benzyladenine (1 mmol) in acetic acid buffer (pH = 4) was treated with bromine (1 mmol) at room temperature for 12 h. The reaction was quenched with 10 % sodium sulfite solution and extracted with dichloromethane. The combined extracts were dried (Na₂SO₄) and evaporated
30 to dryness. Chromatography afforded N⁹-benzyl-8-bromoadenine.
- Step C. A mixture of N⁹-benzyl-8-bromoadenine (1 mmol), aminoethylphosphonate (2 mmol), and sodium hydroxide (2 mmol) in ethanol-water in a sealed tube was warmed at 110 °C under nitrogen. After 24 h the cooled reaction mixture was purified through preparative HPLC to give N⁹-
35 benzyl-8-(2-phosphonoethylamino)adenine (3.1). Exact mass calculated for C₁₄H₁₇N₆O₃P + H⁺: 349.1178. Found: 349.1180.

The following compounds were prepared according to this procedure:

- 3.2: N⁹-phenethyl-8-(2-phosphonoethylamino)adenine. mp 159-160 °C; Anal. calcd. for C₁₅H₁₉N₆O₃P + 1.25 H₂O: C: 46.81; H: 5.63; N: 21.84. Found: C: 47.05; H: 5.63; N: 21.48.
- 3.3: N⁹-(2-naphthylmethyl)-8-(2-phosphonoethylamino)adenine. mp 189-190 °C; Anal. calcd. for C₁₈H₁₉N₆O₃P + 1.5 H₂O: C: 50.82; H: 5.21; N: 19.76. Found: C: 50.71; H: 5.25; N: 19.54.
- 3.4: N⁹-cyclohexylethyl-8-(2-phosphonoethylamino)adenine. mp > 250 °C; Anal. calcd. for C₁₅H₂₅N₆O₃P + 0.33 H₂O: C: 48.12; H: 6.91; N: 22.44. Found: C: 48.12; H: 6.78; N: 22.15.

Example 4.

Preparation of 8-(2-phosphonoethylamino)adenosines.

- 15 A mixture of 8-bromoadenosine (1 mmol), aminoethylphosphonate (2 mmol), and sodium hydroxide (2 mmol) in ethanol-water in a sealed tube was warmed at 110 °C under nitrogen. After 24 h the cooled reaction mixture was purified through preparative HPLC to give 8-(2-phosphonoethylamino)-adenosine (4.1). mp 175 °C; Anal. calcd. for C₁₂H₁₉N₆O₇P + 0.5 H₂O: C: 36.10; H: 5.05; N: 21.05; P: 7.76. Found: C: 36.08; H: 4.83; N: 20.36; P: 7.86.

20 The following compound was prepared in this manner:

4.2: 8-(2-Phosphonoethylamino)-5-deoxyadenosine as a white solid. mp 220 °C; Anal. calcd. for C₁₂H₁₉N₆O₆P + 1.5 H₂O: C: 35.92; H: 5.53; N: 20.94. Found: C: 36.15; H: 5.12; N: 20.53.

25

Examples 5.

Preparation of N⁹-alkyl-8-(phosphonomethoxymethyl)adenines.

- Step A. A mixture of N⁹-phenethyl-8-bromoadenine (1 mmol), tetrakis (triphenylphosphine)palladium (0.05 mmol), and triethylamine (5 mmol) in DMF in a sealed tube was warmed at 110 °C under 50 psi of carbon monoxide. After 24 h the cooled reaction mixture was evaporated and purified through chromatography to gave N⁹-phenethyl-8-methoxycarbonyladenine as a yellow solid. TLC: R_f = 0.12, 5 % MeOH-CH₂Cl₂.
- 30

Step B. A solution of N⁹-phenethyl-8-methoxycarbonyladenine (1 mmol) in tetrahydrofuran was treated with lithium aluminum hydride (1 mmol) at 0 °C for 1 h. Extraction and chromatography gave N⁹-phenethyl-8-hydroxymethyl-adenine as a white solid. TLC: R_f = 0.31, 10 % MeOH-CH₂Cl₂.

- 5 **Step C.** A solution of N⁹-phenethyl-8-hydroxymethyladenine (1 mmol) in methylene chloride was treated with phosphorus tribromide (1 mmol) at 25 °C for 1 h. Extraction and chromatography gave N⁹-phenethyl-8-bromomethyl-adenine as a white solid. TLC: R_f = 0.31, 10 % MeOH-CH₂Cl₂.

- 10 **Step D.** A solution of N⁹-phenethyl-8-bromomethyladenine (1 mmol) in DMF was treated with a solution of diethyl hydroxymethylphosphonate sodium salt (1 mmol) in DMF at 25 °C for 1 h. Extraction and chromatography gave N⁹-phenethyl-8-diethylphosphonomethoxymethyladenine as a white solid. TLC: R_f = 0.31, 10 % MeOH-CH₂Cl₂.

- 15 N⁹-phenethyl-8-diethylphosphonomethoxymethyladenine was subjected to Step F in Example 2 to give N⁹-(2-phenethyl)-8-(phosphonomethoxymethyl)-adenine (5.1) as a white solid. mp > 250 °C; Anal. calcd. for C₁₉H₂₂N₅O₄P + 0.75 H₂O: C: 56.93; H: 5.91; N: 10.48. Found: C: 56.97; H: 5.63; N: 10.28.

The following compounds were prepared according to this procedure:

- 20 **5.2:** N⁹-(2-cyclohexylethyl)-8-(phosphonomethoxymethyl)adenine. mp > 250 °C; Anal. calcd. for C₁₅H₂₄N₅O₄P + 1 H₂O: C: 46.51; H: 6.76; N: 18.08. Found: C: 46.47; H: 6.71; N: 17.91.
- 5.3:** N⁹-(1-nanonyl)-8-(phosphonomethoxymethyl)adenine. mp 195-210 °C; Anal. calcd. for C₁₆H₂₈N₅O₄P + 1 H₂O: C: 47.64; H: 7.50; N: 17.36. Found: C: 25 47.33; H: 7.34; N: 16.99.
- 5.4:** N⁹-(3-cyclohexylpropyl)-8-(phosphonomethoxymethyl)adenine. mp 230-250 °C; Anal. calcd. for C₁₉H₂₂N₅O₄P + 0.9 H₂O + 0.3 HBr: C: 45.34; H: 6.68; N: 16.52. Found: C: 45.74; H: 6.39; N: 16.18.

- 30 Alternatively this type of compound can also be prepared according to the following procedure:

- Step E.** A solution of 6-chloro-5-amino-4-(neopentylamino)pyrimidine (1 mmol) in diethyl ether was treated with pyridine (3 mmol), and acetoxyacetyl chloride (1.2 mmol) at 25 °C for 12 h. Extraction and chromatography afforded 35 6-chloro-5-acetoxyacetyl-amino-4-neopentylaminopyrimidine as a yellow solid. TLC: R_f = 0.18, 30 % EtOAc-hexane.

Step F. A solution of 6-chloro-5-acetoxyac tylamino-4-neopentyl-aminopyrimidine (1 mmol) in phosphorus oxychloride was heated at reflux for 6 h. The cooled reaction mixture was evaporated to dryness and the residue was dissolved in pyridine and stirred at 25 °C for 20 h. Evaporation and chromatography afforded 6-chloro-8-acetoxymethyl-N⁹-neopentylpurine as a yellow solid. TLC: R_f = 0.51, 50 % EtOAc-hexane.

Step G. A solution of 6-chloro-8-acetoxymethyl-N⁹-neopentylpurine (1 mmol) in THF-water was treated with aqueous sodium hydroxide (1.5 mmol) at 0 °C for 0.5 h. Extraction and chromatography afforded 6-chloro-8-hydroxymethyl-N⁹-neopentylpurine as a yellow gel. TLC: R_f = 0.38, 33 % EtOAc-hexane.

Step H. A solution of 6-chloro-8-hydroxymethyl-N⁹-neopentylpurine (1 mmol) in methylene chloride was treated with phosphorus tribromide (1 mmol) at 25 °C for 6 h. Extraction and chromatography afforded 6-chloro-8-bromomethyl-N⁹-neopentylpurine as a white solid. TLC: R_f = 0.64, 25 % EtOAc-hexane.

Step I. A solution of 6-chloro-8-bromomethyl-N⁹-neopentylpurine (1 mmol) in DMF was treated with a solution of sodium diethylphosphonmethoxide (1 mmol) at 25 °C for 6 h. Extraction and chromatography afforded 6-chloro-8-diethyl-phosphonmethoxymethyl-N⁹-neopentylpurine as a white solid. TLC: R_f = 0.31, 50 % EtOAc-hexane.

Step J. A solution of 6-chloro-8-diethylphosphonmethoxymethyl-N⁹-neopentylpurine (1 mmol) in THF-DMSO was treated with liquid ammonia (10 mmol) at 25 °C for 6 h. Extraction and chromatography afforded 8-diethylphosphonmethoxymethyl-N⁹-neopentyladenine as a white solid. TLC: R_f = 0.44, 25 % MeOH-EtOAc.

8-Diethylphosphonmethoxymethyl-N⁹-neopentyladenine was subjected to Step F in Example 2 to give N⁹-neopentyl-8-(phosphonmethoxymethyl)-adenine (5.5) as a white solid. mp > 250 °C; Anal. calcd. for C₁₂H₂₀N₅O₄P + 1.5 H₂O: C: 40.45; H: 6.51; N: 19.65. Found: C: 40.68; H: 6.35; N: 19.40.

Examples 6.

Preparation of N⁹-substituted-8-(1-(3-phosphono)propyl)adenines.

Step A. A mixture of diethyl propargylphosphonate (1 mmol, prepared according to *J. Org. Chem.*, 1993, 58(24), 6531.), tributyltin hydride (1.05 mmol), and AIBN (0.005 mmol) was heated at 60 °C for 18 h. The cooled

reaction mixture was purified through chromatography to give dimethyl (1-tributylstannyl)allyl-3-phosphonate as a yellow oil.

Step B. A solution of N⁹-(2-cyclohexylethyl)-8-bromoadenine (1 mmol), tetrakis(triphenylphosphine)palladium (0.1 mmol), and dimethyl (1-tributylstannyl)allyl-3-phosphonate (5 mmol) in DMF was warmed at 90 °C under nitrogen. After 2 h the cooled reaction mixture was evaporated and purified through chromatography to give N⁹-(2-cyclohexylethyl)-8-(3-dimethylphosphonopropene-1-yl)adenine as a yellow solid. TLC: R_f = 0.48, 10 % MeOH - CH₂Cl₂.

Step C. A solution of N⁹-(2-cyclohexylethyl)-8-(3-dimethylphosphonopropene-1-yl)adenine in methanol-acetic acid was stirred at room temperature under 50 psi of H₂ for 12 h. Filtration and chromatography afforded N⁹-(2-cyclohexylethyl)-8-(1-(3-dimethylphosphono)propyl)adenine as a yellow solid. TLC: R_f = 0.26, 10 % MeOH - CH₂Cl₂.

N⁹-(2-cyclohexylethyl)-8-(1-(3-dimethylphosphono)propyl)adenine was subjected to Step F in Example 2 to give N⁹-(2-cyclohexylethyl)-8-(1-(3-phosphono)propyl)adenine (6.1) as a white solid: mp 122-125 °C; Anal. calcd. for C₁₆H₂₆N₅O₃P + 0.25 AcOH: C: 51.83; H: 7.12; N: 18.31. Found: C: 51.87; H: 6.96; N: 17.96.

6.2: N⁹-(2-phenethyl)-8-(1-(3-phosphono)propyl)adenine was also prepared in this manner as a solid. mp > 250 °C Anal. calcd. for C₁₆H₂₀N₅O₃P + 0.5 H₂O: C: 51.89; H: 5.71; N: 18.91. Found: C: 51.81; H: 5.49; N: 18.66.

Examples 7.

Preparation of N⁹-(2-phenethyl)-8-(2-(5-phosphono)thienyl)adenine.

Step A. A solution of 2-thienyllithium in THF (1 mmol) was added to a solution of diethyl chlorophosphate (1 mmol) at -78 °C under nitrogen. After 2 h the reaction was warmed to room temperature and quenched with brine. Extraction and chromatography afforded 2-diethylphosphonothiophene as a yellow oil. TLC: R_f = 0.37, 50 % EtOAc - hexane.

Step B. A solution of 2-diethylphosphonothiophene (1 mmol) in THF was treated with nBuLi at -78 °C for 1 h. Tributyltin chloride was added and stirred at -78 °C for 2 h and the reaction was quenched with water and warmed to room temperature. Extraction and chromatography afforded diethyl 2-(5-tributylstannyl)thienylphosphonate as a yellow oil. TLC: R_f = 0.65, 50 % EtOAc - hexane.

Step C. A mixture of N⁹-phenethyl-8-bromoadenine (1 mmol), tetrakis (triphenylphosphine)palladium (0.1 mmol), and diethyl 2-(5-tributylstannyl)-thienylphosphonate (5 mmol) in DMF was warmed at 80 °C under nitrogen. After 21 h the cooled reaction mixture was evaporated to dryness. The dark oil
5 was triturated with hexane and the residue was dissolved in CH₂Cl₂ and filtered. The filtrate was evaporated to give N⁹-(2-phenethyl)-8-(2-(5-diethylphosphono)-thienyl)adenine as a yellow solid. TLC: R_f = 0.50, 10 % MeOH - CH₂Cl₂.

N⁹-(2-phenethyl)-8-(2-(5-diethylphosphono)thienyl)adenine was subjected to Step F in Example 2 to give N⁹-(2-phenethyl)-8-(2-(5-phosphono)-thienyl)adenine (7.1) as a white solid. mp > 250 °C; Anal. calcd. for
10 C₁₇H₁₆N₅O₃SP + 0.5H₂O: C: 49.76; H: 4.17; N: 17.07. Found: C: 50.07; H: 4.02; N: 17.45.

N⁹-(2-phenethyl)-8-(2-(5-phosphono)thienyl)adenine can also be made via a cyclization reaction between 5-diethylphosphono-2-thiophene-
15 carboxaldehyde (prepared from 2-thienyllithium as described in Steps C and D of Example 1) as described in Example 2.

Example 8.

Preparation of N⁹-(2-cyclohexylethyl)-8-(phosphonomethylthio)-adenine.

Step A. A mixture of N⁹-(2-cyclohexylethyl)-8-bromoadenine (1 mmol), and K₂S (4 mmol) in ethanol was warmed at 110 °C for 7 h, and at 85 °C for 12 h. The cooled reaction mixture was filtered, evaporated and purified through
20 chromatography to give N⁹-(2-cyclohexylethyl)-8-thiohydroxyadenine as a yellow solid. TLC: R_f = 0.26, 5 % MeOH - CH₂Cl₂.

Step B. A mixture of N⁹-(2-cyclohexylethyl)-8-thiohydroxyadenine (1 mmol), K₂CO₃ (4 mmol), and diethyl chloromethylphosphonate (3 mmol) in DMF was stirred at room temperature for 48 h. Extraction and chromatography gave
25 N⁹-(2-cyclohexylethyl)-8-diethylphosphono-methylthioadenine. TLC: R_f = 0.35, 10 % MeOH - EtOAc.

N⁹-(2-Cyclohexylethyl)-8-diethylphosphonomethylthioadenine was subjected to Step F in Example 2 to give N⁹-(2-cyclohexylethyl)-8-(phosphonomethylthio)adenine (8.1) as a white solid. mp 240 - 243 °C; Anal.
30 Calcd. for C₁₄H₂₂N₅O₃SP + 1.25H₂O: C: 42.69; H: 5.95; N: 17.54. Found: C: 42.62; H: 6.03; N: 17.80.

35

Example 9.**Preparation of 6-chloro-9-phenethyl-8-(2-(5-phosphono)furanyl)purine.**

6-Chloro-N⁹-phenethyl-8-(2-(5-diethylphosphono)furanyl)purine (Step C in Example 2) was subjected to procedure of Step F in Example 2 to give
5 compound 9.1 as a yellow solid. mp >200 °C; Anal. calcd. for C₁₇H₁₄N₄O₄PCl + 2 H₂O + 0.28 HBr: C: 44.06; H: 3.98; N: 12.09. Found: C: 43.86; H: 3.59; N: 12.02.

Example 10.**Preparation of N⁶,N⁹-substituted-8-(2-(5-phosphono)furanyl)adenines.**

10 A solution of 6-chloro-N⁹-substituted-8-(2-(5-diethylphosphono)furanyl)-purine (1 mmol) in DMSO was treated with alkylamine at 100 °C for 12 h. Evaporation and chromatography gave N⁶,N⁹-substituted-8-(2-(5-diethylphosphono)furanyl)adenines.

The title compounds were obtained by subjecting N⁶,N⁹-substituted-8-(2-(5-diethylphosphono)furanyl)adenines to the procedure of Step F in Example 2.
15

The following compounds were prepared in this manner:

10.1: 6-Dimethylamino-N⁹-phenethyl-8-(2-(5-phosphono)furanyl)purine as a white solid. mp >200 °C; Anal. calcd. for C₁₉H₂₀N₅O₄P: C: 55.2; H: 4.8; N: 16.9. Found: C: 54.9; H: 4.9; N: 16.6.

20 **10.2:** 6-Methylamino-N⁹-phenethyl-8-(2-(5-phosphono)furanyl)purine as a white solid. mp 242 °C; Anal. calcd. for C₁₈H₁₈N₅O₄P + 1 H₂O: C: 51.8; H: 4.8; N: 16.8. Found: C: 51.7; H: 4.8; N: 16.7.

Example 11.

25 **Preparation of 2-methylthio-6-amino-N⁹-isobutyl-8-(2-(5-phosphono)furanyl)purine and 2-methylsulfonyl-6-amino-N⁹-isobutyl-8-(2-(5-phosphono)furanyl)purine.**

Step A: 2-Methylthio-4,5,6-triaminopyrimidine and 5-diethylphosphono-2-furaldehyde was subjected to the procedures of Step D in Example 2 to give 6-amino-2-methylthio-8-(2-(5-diethylphosphono)furanyl)purine as a yellow solid.
30 TLC: R_f = 0.27, 80 % EtOAc - hexane.

Step B: 6-Amino-2-methylthio-8-(2-(5-diethylphosphono)furanyl)purine was alkylated with isobutyl bromide following the procedures of Step A in Example 3 to give 6-amino-N⁹-isobutyl-2-methylthio-8-(2-(5-diethylphosphono)-
35 furanyl)purine as a yellow solid. TLC: R_f = 0.27, 80 % EtOAc - hexane.

- Step C:** 6-Amino-N⁹-isobutyl-2-methylthio-8-(2-(5-diethyl-phosphono)-furanyl)purine was subjected to Step F in Example 2 to give 6-amino-N⁹-isobutyl-2-methylthio-8-(2-(5-phosphono)-furanyl)purine (11.1) as a white solid. mp 220 °C; Anal. calcd. for C₁₄H₁₈N₅O₄PS + 0.25 HBr + 0.25 EtOAc: C: 42.33; H: 4.8; N: 16.45. Found: C: 42.42; H: 4.53; N: 16.39.
- Step D:** A solution of 6-amino-N⁹-isobutyl-2-methylthio-8-(2-(5-diethyl-phosphono)furanyl)purine (1 mmol) in 50 mL of methanol was cooled to 0 °C and treated with an acetone solution of Oxone (1.6 mmol). After stirring for 3 h at 25 °C the reaction was extracted and then chromatographed to give 6-amino-N⁹-isobutyl-2-methylsulfonyl-8-(2-(5-diethylphosphono)furanyl)purine as a white solid. TLC: R_f = 0.24, 100 % EtOAc.
- Step E:** 6-Amino-N⁹-isobutyl-2-methylsulfonyl-8-(2-(5-diethylphosphono)-furanyl)purine was subjected to Step F in Example 2 to give 6-amino-N⁹-isobutyl-2-methylsulfonyl-8-(2-(5-phosphono)furanyl)purine (11.2) as a white solid. mp 240 °C (decomp); Anal. calcd. for C₁₄H₁₈N₅O₆PS + 0.5 H₂O: C: 39.62; H: 4.51; N: 16.5. Found: C: 39.77; H: 4.44; N: 16.12.

Example 12.

Preparation of 6-amino-N⁹-neopentyl-8-(2-(3,4-dichloro-5-phosphono)furanyl)purine.

- Step A:** A solution of 3,4-dichloro-2-furoic acid (1 mmol) in diethyl ether was treated with LDA (3 mmol) at -78 °C for 30 min and then treated with diethyl chlorophosphate (3.5 mmol) at -78 °C for 1 h. The reaction was quenched and extracted to give 5-diethylphosphono-3,4-dichloro-2-furoic acid as a yellow foam.
- Step B:** A solution of 5-diethylphosphono-3,4-dichloro-2-furoic acid (1 mmol) in methylene chloride was treated with oxalyl chloride and DMF at 25 °C for 1 h. The reaction mixture was evaporated and the residue was dissolved in diethyl ether and treated with a solution of 4-chloro-5-amino-6-neopentylaminopyrimidine (1 mmol) and pyridine (3 mmol) in diethyl ether at 25 °C for 16 h. Extraction and chromatography gave 4-chloro-5-(2-(3,4-dichloro-5-diethylphosphono)furoyl)amino-6-neopentylaminopyrimidine as a yellow solid. TLC: R_f = 0.4, 50 % EtOAc - hexane.
- Step C:** A solution of 4-chloro-5-(2-(3,4-dichloro-5-diethylphosphono)-furoyl)amino-6-neopentylaminopyrimidine (1 mmol) in dichloromethane was treated with silicone tetrachloride (2.5 mmol) and triethylamine (2.5 mmol) at 45

°C for 18 h. The cooled reaction mixture was subjected to extraction and chromatography to give 6-chloro-N⁹-neopentyl-8-(2-(3,4-dichloro-5-diethylphosphono)furanyl)purine as a yellow solid. TLC: R_f = 0.28, 50 % EtOAc - hexane.

- 5 Step D: 6-Chloro-N⁹-neopentyl-8-(2-(3,4-dichloro-5-diethylphosphono)-furanyl)purine was subjected to Steps E and F in Example 2 to give 6-amino-N⁹-neopentyl-8-(2-(3,4-dichloro-5-phosphono)furanyl)purine (**12.1**) as a white solid. mp > 250 °C; Anal. calcd. for C₁₄H₁₆N₅O₄PCl₂ + 0.5 H₂O + 0.15 EtOAc: C: 39.64; H: 4.15; N: 15.83. Found: C: 39.82; H: 3.88; N: 15.46.

10

Example 13.

Preparation of hydroxyethyl disulfidylethylphosphonate diester.

- A suspension of 8-(2-(5-phosphono)furanyl)-N⁹-phenethyladenine (1 mmol) in thionyl chloride (5 mL) was warmed at reflux for 4 h. The cooled
15 reaction mixture was evaporated to dryness and the resulting yellow residue was treated with a solution of 2-hydroxyethyl disulfide (4 mmol), pyridine (2.5 mmol) in methylene chloride. After stirring at 25 °C for 4 h the reaction was subjected to extraction and chromatography to give two compounds:
20 **13.1:** N⁹-phenethyl-8-(bis(6'-hydroxy-3',4'-disulfide)hexylphosphono)furanyl)-adenine. Anal. calcd for C₂₅H₃₂N₅O₆S₄P + 0.5 DMSO + 1.5 H₂O: C: 43.15; H: 5.29; N: 9.68. Found: C: 43.38; H: 4.93; N: 9.34.
13.2: N⁹-phenethyl-8-((3',4'-disulfide)nonacyclicphosphono)furanyl)-adenine. Anal. calcd for C₂₁H₂₂N₅O₄S₂P + DMSO: C: 47.49; H: 4.85; N: 12.04. Found: C: 47.93; H: 4.60; N: 11.76.

25

Example 14.

Preparation of substituted benzyl phosphonate diesters.

- A suspension of 8-(2-(5-phosphono)furanyl)-N⁹-phenethyladenine (1 mmol) in thionyl chloride (5 mL) was warmed at refluxing 4 h. The cooled
30 reaction mixture was evaporated to dryness and a solution of the resulting yellow residue was added to a solution of the corresponding benzyl alcohol (4 mmol), and pyridine (2.5 mmol) in methylene chloride. After stirring at 25 °C for 4 h the reaction mixture was subjected to extraction and chromatography to give the title compounds.

14.1: N⁹-phenethyl-8-(2-(5-(bis-(3-bromo-4-methoxy)benzyl)phosphono)-furanyl)adenine. Molecular mass calculated for C₃₃H₃₀N₅O₆Br₂P + H⁺: 784.

Found: 784.

14.2: N⁹-phenethyl-8-(2-(5-(bis-(3-cyano-4-methoxy)benzyl)phosphono)-furanyl)adenine. Anal. calcd. for C₃₅H₃₀N₇O₆P + 0.5 H₂O : C: 61.40; H: 4.56; N: 14.32. Found: C: 61.45; H: 4.51; N: 14.18.

14.3: N⁹-neopentyl-8-(2-(5-(bis-(4-acetoxy)benzyl)phosphono)furanyl)adenine. Anal. calcd. for C₃₂H₃₄N₅O₈P + 0.6 H₂O : C: 58.37; H: 5.39; N: 10.64. Found: C: 58.11; H: 5.28; N: 10.42.

10 N⁹-neopentyl-8-(2-(5-(bis-(3-phthalidyl-2-ethyl)phosphono)furanyl)-adenine is also prepared following the above described procedure using 2-(3-phthalidyl)ethanol which was prepared from phthalide-3-acetic acid in Example 27.

This reaction procedure can also be used to prepare diaryl ester
15 prodrugs of phosphonates, such as substituted phenyl esters of phosphonate.

Example 15.

Preparation of 6-amino-8-(2-(5-diphenylphosphono)furanyl)-N⁹-(2-phenyl)ethylpurine.

20 **Step A.** A suspension of 6-chloro-8-(2-furanyl)-N⁹-phenethylpurine (1 mmol) in THF at -78 °C was treated with LDA (1.3 mmol) for 1 h. Then a solution of diphenyl chlorophosphate in THF was added and the reaction was stirred at -78 °C for another hour. The reaction was warmed to 0 °C and quenched with aqueous saturated sodium bicarbonate. Extraction and chromatography gave
25 6-chloro-8-(2-(5-diphenylphosphono)furanyl)-N⁹-phenethylpurine as a white solid. mp 117-118 °C.

Step B. A solution of 6-chloro-8-(2-(5-diphenylphosphono)furanyl)-N⁹-phenethylpurine (1 mmol) in DMF was treated with sodium azide (4 mmol) and triphenylphosphine (4 mmol) at room temperature for 3 h. Filtration, evaporation
30 of the filtrate followed by chromatography gave 6-triphenyl-phosphonoimino-8-(2-(5-diphenylphosphono)furanyl)-N⁹-(2-phenyl)ethylpurine as a beige foam.

Step C. A solution of 6-triphenylphosphonoimino-8-(2-(5-diphenylphosphono)furanyl)-9-(2-phenyl)ethylpurine (1 mmol) in THF was treated with aqueous hydrogen chloride at room temperature for 24 h. Evaporation and chromatography gave 6-amino-8-(2-(5-diphenylphosphono)furanyl)-N⁹-(2-phenyl)ethylpurine (**15.1**) as a pale yellow solid. mp 196-197 °C; Anal. calcd.
35

for $C_{29}H_{24}N_5O_4P$: C: 64.80; H: 4.50; N: 13.03; P: 5.76. Found: C: 64.50; H: 4.47; N: 12.98; P: 5.46.

Example 16.

5 Preparation of acyloxymethylphosphonate diesters.

A solution of 8-(2-(5-phosphono)furanyl)- N^9 -phenethyladenine (1 mmol) in acetonitrile and N,N,N-diisopropylethylamine (5 mmol) was treated with acyloxymethyl iodide (4 mmol) at 0 °C for 24 h. Extraction and chromatography gave the title compounds.

10 The following compounds were prepared according to this procedure:

16.1: 6-Amino-9-phenethyl-8-(2-(5-diisobutyrylmethylphosphono)furanyl)-purine. Anal. calcd for $C_{27}H_{32}N_5O_8P$: C: 55.40; H: 5.50; N: 12.00. Found: C: 55.60; H: 5.60; N: 11.80.

15 16.2: 6-Amino-9-(2-cyclohexylethyl)-8-(2-(5-diisobutyrylmethylphosphono)furanyl)purine. Anal. calcd for $C_{27}H_{38}N_5O_8P + 0.7 H_2O$: C: 53.70; H: 6.60; N: 11.60. Found: C: 54.00; H: 6.50; N: 11.20.

16.3: 6-Amino-9-ethyl-8-(2-(5-diisobutyrylmethylphosphono)furanyl)purine. Anal. calcd for $C_{21}H_{28}N_5O_8P$: C: 49.51; H: 5.54; N: 13.75. Found: C: 49.75; H: 5.37; N: 13.76.

20 16.4: 6-Amino-9-neopentyl-8-(2-(5-diisobutyrylmethylphosphono)furanyl)-purine. Anal. calcd for $C_{24}H_{34}N_5O_8P$: C: 52.27; H: 6.21; N: 12.70. Found: C: 52.40; H: 6.27; N: 12.41.

16.5: 6-Amino-9-neopentyl-8-(2-(5-dipivaloxymethylphosphono)furanyl)purine. Anal. calcd for $C_{26}H_{38}N_5O_8P + 0.2 EtOAc$: C: 53.90; H: 6.68; N: 11.73. Found: C:

25 54.10; H: 6.80; N: 11.42.

6-Amino-9-phenethyl-8-(2-(5-bis-(3-(5,6,7-trimethoxy)phthalidyl)-phosphono)furanyl)purine (**16.6**) was also synthesized following this procedure using 3-bromo-5,6,7-trimethoxyphthalide as the alkylating reagent to give the titled compound as a white solid after preparative HPLC purification.

30 mp 155-160 °C; Anal. calcd. for $C_{39}H_{36}N_5O_{14}P + H_2O$: C: 55.26; H: 4.52; N: 8.26. Found: C: 54.89; H: 4.75; N: 8.21.

Example 17.**Preparation of 5-methyl-4-hydroxymethyl-2-oxo-1,3-dioxolene.**

A solution of 4,5-dimethyl-2-oxo-1,3-dioxolene (1 mmol) and selenium dioxide (2.5 mmol) in dioxane was heated at reflux for 1 h. Evaporation,
5 extraction and chromatography gave 5-methyl-4-hydroxymethyl-2-oxo-1,3-dioxolene as a yellow oil. TLC: $R_f = 0.5$, 5 % MeOH-dichloromethane.

Example 18.**Preparation of (5-substituted 2-oxo-1,3-dioxolen-4-yl)methyl phosphonate**
10 **prodrugs.**

A solution of N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine (1 mmol) in DMF and 2 mmol of sodium hydride is treated with 5-methyl-4-bromomethyl-2-oxo-1,3-dioxolene (4 mmol, prepared according to *Chem. Pharm. Bull.* 1984, 32(6), 2241) at 25 °C for 24 h. Extraction and chromatography gives N⁹-
15 neopentyl-8-(2-(5-bis(5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl phosphono)-furanyl)adenine.

Alternatively, N⁹-neopentyl-8-(2-(5-bis(5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl phosphono)-furanyl)adenine is prepared from N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine and 5-methyl-4-hydroxymethyl-2-oxo-1,3-dioxolene (prepared from 4,5-dimethyl-2-oxo-1,3-dioxolene as described in
20 Example 17) according to procedures of Example 14.

Example 19.**Preparation of 2-(6-amino-N⁹-neopentylpurin-8-yl)phenyl phosphonate:**

25 **Step A.** Triethylamine (1.1 mmol) was added slowly to an ice-cooled solution of 6-chloro-N⁹-neopentyl-8-(2-hydroxyphenyl)purine (1 mmol) and diethyl phosphite (1 mmol) in carbontetrachloride. The reaction was stirred at room temperature overnight. Triethylamine hydrochloride was precipitated as a white solid mass. Extraction and chromatography gave diethyl 2-(6-Chloro-N⁹-neopentylpurin-8-yl)phenyl phosphate.
30

Step B. Diethyl 2-(6-Chloro-N⁹-neopentylpurin-8-yl)phenyl phosphate was subjected to Step E and F in Example 2 to give the title compound (19.1). mp > 250° C; Anal. calcd. for C₁₆H₂₀N₅O₄P + 1.25 H₂O: C: 48.06; H: 5.67; N: 17.51. Found: C:48.42; H: 5.42; N: 17.15.

35

Example 20.Preparation of N⁹-neopentyl-8-(1-(2-phosphono)imidazolemethyl)adenine:

Step A. A solution of 1-benzylimidazole (1.1 mmol) in THF was treated with LDA (1.1 mmol) at -78 °C for 1 h, and followed by addition of diethyl chlorophosphate (2 mmol), and stirred for 2 h. Extraction and chromatography gave 1-benzyl-2-diethylphosphonoimidazole as a yellow oil. TLC: R_f = 0.35, 80 % EtOAc-hexane.

Step B. A solution of 1-benzyl-2-diethylphosphonoimidazole (1 mmol) in EtOH was treated with palladium on carbon (10 %) at 25 °C under 1 atmosphere of hydrogen for 19 h. Filtration and evaporation gave 2-diethylphosphonoimidazole as a white solid. TLC: R_f = 0.05, 80 % EtOAc-hexane.

Step C. A solution of 8-bromomethyl-6-chloro-N⁹-neopentylpurine (1 mmol, Step H of Example 5), 2-diethylphosphonoimidazole (2.5 mmol), and N,N,N-diisopropylethylamine (2.5 mmol) in acetonitrile was stirred at 25 °C for 48 h. Extraction and chromatography gave 6-chloro-N⁹-neopentyl-8-(1-(2-diethylphosphono)imidazolemethyl)purine.

Step D. 6-Chloro-N⁹-neopentyl-8-(1-(2-diethylphosphono)imidazolemethyl)purine was subjected to Steps E and F in Example 2 to give the title compound (20.1). mp > 250 °C; MS (M + H) calcd. for C₁₄H₂₀N₇O₃P: 366; found: 366.

Example 21.Preparation of N⁹-phenethyl-8-(phosphonomethylaminocarbonyl)adenine:

Step A. N⁹-phenethyl-8-(methoxycarbonyl)adenine (1 mmol, prepared as in Step A of Example 5) was treated with sodium hydroxide (1.2 mmol) in THF:MeOH:H₂O (3:2:1) at 25 °C for 1.5 h. The reaction mixture was evaporated to dryness, and the residue was dissolved in DMF, treated with diethyl aminomethylphosphonate (1.5 mmol), EDCI (1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide hydrochloride, 1.3 mmol), HOBt (1-hydroxy-benzotriazole hydrate, 1.5 mmol), and stirred at 25 °C for 24 h. Extraction and chromatography gave N⁹-phenethyl-8-(diethylphosphonomethylaminocarbonyl)adenine as a white solid. TLC: R_f = 0.1, EtOAc.

Step B. N⁹-phenethyl-8-(diethylphosphonomethylaminocarbonyl)adenine (1 mmol) was subjected to Step F in Example 2 to give the title compound (21.1). mp > 250 °C; Anal. calcd. for C₁₅H₁₇N₆O₄P + 0.17 Toluene: C: 47.32; H: 4.50; N: 20.45. Found: C:47.67; H: 4.57; N: 20.78.

Example 22.Preparation of 2-substituted N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine:

Step A. A solution of 2-amino-4,6-dichloropyrimidine (1 mmol),
5 neopentylamine (1.05 mmol), and triethylamine (2 mmol) in n-butanol was stirred at 110 °C for 12 h. Extraction and chromatography gave 2-amino-4-chloro-6-neopentylpyrimidine as a yellow solid. TLC: R_f = 0.2, 30 % EtOAc-hexane.

Step B. A mixture of 2-amino-4-chloro-6-neopentylpyrimidine (1 mmol),
10 sodium acetate (14 mmol), acetic acid (86 mmol), and 4-chlorobenzene-diazonium hexafluorophosphate (1.15 mmol) in water was stirred at 25 °C for 12 h. Extraction and evaporation gave a yellow solid which was treated with zinc dust (10 mmol) and acetic acid (5.54 mmol) in EtOH-H₂O at 80 °C for 1 h. Extraction and chromatography gave 4-chloro-2,5-diamino-6-neopentylpyrimidine as a yellow solid. TLC: R_f = 0.25, 50 % EtOAc-hexane.

Step C. 4-Chloro-2,5-diamino-6-neopentylpyrimidine was subjected to
Step D, E, F in Example 2 to give 2,6-diamino-N⁹-neopentyl-8-(2-(5-phosphono)furanyl)purine (22.1) as a yellow solid. mp 240° C (decomp); Anal. calcd. for C₁₄H₁₉N₆O₄P + 2.2 HBr + 0.5 acetone: C: 32.47; H: 4.25; N: 14.66.
20 Found: C:32.31; H: 4.51; N: 14.85.

Similarly, 2-methylthio-N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine (22.2) was also prepared from 4-amino-6-chloro-2-methylthiopyrimidine as a yellow solid. mp > 250; Anal. calcd. for C₁₅H₂₀N₅O₄PS + 0.2 CH₂Cl₂ + 0.1 toluene: C: 45.08; H: 5.04; N: 16.53. Found: C:45.27; H: 5.34; N: 16.24.

Example 23.Preparation of alkyloxycarbonyloxyalkyl phosphonate esters.

A solution of N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine (1 mmol) in 5 mL of anhydrous DMF is treated with N,N'-dicyclohexyl-4-morpholinecarboxamide (5 mmol), and isopropylloxycarbonyloxymethyl iodide (5 mmol) which is prepared from the commercially available
30 chloromethyl chloroformate according to the reported procedure, Nishimura et al. *J. Antibiotics*, 1987, 40(1), 81-90. The reaction mixture is stirred for 24 h at room temperature and the solvent is removed under reduced pressure. The resulting syrup is chromatographed on silica with 50% /50% EtOAc/Hexane to
35 yield N⁹-neopentyl-8-(2-(5-diisopropylloxycarbonyloxymethyl phosphono)furanyl)adenine.

Other alkyloxycarbonyloxymethyl, aryloxycarbonyloxymethyl, alkyl- and arylthiocarbonyloxymethyl phosphonate esters can also be prepared following the above described procedure.

5 Example 24.

Preparation of 1-substituted-1,3-propanediol cyclic esters of purine phosphonates.

Step A(J. Org. Chem., 1957, 22, 589)

To a solution of 2-pyridine propanol (72.9 mmol) in acetic acid (75 mL) was
10 added 30% hydrogen peroxide slowly. The reaction mixture was heated to 80 °C for 16 h. The reaction was concentrated under vacuum and the residue was dissolved in acetic anhydride (100 mL) and heated at 110 °C overnight. Acetic anhydride was evaporated upon completion of reaction. Chromatography of the mixture by eluting with methanol-methylene chloride (1:9) resulted in 10.5 g
15 of pure 2-(1-(1,3-diacetoxy)propyl)pyridine.

Step B. To a solution of 2-(1-(1,3-diacetoxy)propyl)pyridine (21.1 mmol) in methanol-water (3:1, 40 mL) was added potassium carbonate (105.5 mmol). After stirring for 3 h at room temperature, the reaction mixture was concentrated.
20 The residue was chromatographed by eluting with methanol-methylene chloride (1:9) to give 2.2 g of crystalline 2-(1-(1,3-dihydroxy)propyl)pyridine.

Step C. A suspension of N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine (1 mmol) in 5 mL of thionyl chloride was heated at reflux temperature for 4 h.
25 The reaction mixture was cooled and evaporated to dryness. To the resulting residue was added a solution of 2-(1-(1,3-dihydroxy)propyl)pyridine (1 mmol) and pyridine (2.5 mmol) in 3 mL of methylene chloride. After stirring at 25 °C for 4 h the reaction was subjected to work up and chromatography to give N⁹-neopentyl-8-(2-(5-(1-(2-pyridyl)propan-1,3-yl)phosphono)furanyl)adenine (24.1) as a sticky solid. Anal. Calcd. for C₂₂H₂₅N₈O₄P + 0.75 H₂O + 1.0 HCl: C:50.97; H: 5.35; N: 16.21. Found: C:51.19, H: 5.02; N: 15.91.
30

Following the above described procedures, other cyclic esters are also prepared, such as N⁹-neopentyl-8-(2-(5-(1-(4-pyridyl)propan-1,3-yl)phosphono)furanyl)adenine, N⁹-neopentyl-8-(2-(5-(1-(3-pyridyl)propan-1,3-

yl)phosphono)furanyl)adenine, and N⁹-neopentyl-8-(2-(5-(1-phenylpropan-1,3-yl)phosphono)furanyl)adenine.

Example 25.

5 Preparation of 2-substituted-1,3-propanediol cyclic esters of purine phosphonates.

Step A. To a solution of 2-(hydroxymethyl)-1,3-propanediol (1 g, 9.4 mmol) in pyridine (7.5 mL) at 0 °C was added acetic anhydride (0.89 mL, 9.4 mmol) slowly. The resulting solution was warmed to room temperature and stirred for
10 16 h. The reaction was concentrated under reduced pressure and chromatographed by eluting with methanol-dichloromethane (1:9) to give 510 mg of pure 2-acetoxymethyl-1,3-propanediol.

Step B. 2-Acetoxymethyl-1,3-propanediol was coupled to N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine following Step C of Example 24 to give N⁹-
15 neopentyl-8-(2-(5-(2-(acetoxymethyl)propan-1,3 -yl)phosphono)furanyl)adenine (25.1). mp = 164 - 165 °C; Anal. Calcd. for C₂₀H₂₆N₅O₆P: C: 51.84; H: 5.65; N: 15.11 . Found: C: 52.12; H: 5.77 ; N: 14.59.

Following the above described procedures, other cyclic esters are also prepared, such as N⁹-neopentyl-8-(2-(5-(2-(methoxycarbonyloxymethyl)-propan-1,3 -yl)phosphono)furanyl)adenine, N⁹-neopentyl-8-(2-(5-(2-(hydroxymethyl)-propan-1,3 -yl)phosphono)furanyl)adenine, N⁹-neopentyl-8-(2-(5-(2,2-dihydroxymethylpropan-1,3 -yl)phosphono)furanyl)adenine. N⁹-neopentyl-8-(2-(5-(2-(methoxycarbonyloxymethyl)propan-1,3 -yl)phosphono)-furanyl)adenine is prepared by coupling N⁹-neopentyl-8-(2-(5-phosphono)-
20 furanyl)adenine with 2-(methoxycarbonyloxymethyl)-1,3-propanediol which was prepared as follows:

To a solution of 2-(hydroxymethyl)-1,3-propanediol (9.4 mmol) in dichloromethane (20 mL) and pyridine (7.5 mL) at 0 °C was added methyl chloroformate (9.4 mmol) slowly. The resulting solution was warmed to room
30 temperature and stirred for 16 h. The reaction was concentrated under reduced pressure and chromatographed by eluting with methanol-dichloromethane (1:4) to give 650 mg of 2-(methoxycarbonyloxymethyl)-1,3-propanediol.

Example 26.**Preparation of 8-(2-(5-hydroxyl-1,3 cyclohexyl)phosphono)furanylpurines.**

A suspension of N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine (1 mmol) in 5 mL of thionyl chloride was heated at reflux temperature for 4 h. The reaction mixture was cooled and evaporated to dryness. To the resulting residue was added a solution of cis,cis-1,3,5-cyclohexanetriol (1 mmol) and pyridine (2.5 mmol) in 3 mL of methylene chloride. After stirring at 25 °C for 24 h the reaction was subjected to work up and chromatography to give N⁹-neopentyl-8-(2-(5-(5-hydroxyl-1,3 cyclohexyl)phosphono)furanyl)adenine, minor isomer (26.1). mp 248 - 250 °C; Anal. Cald. for C₂₀H₂₆N₅O₅P + 0.5 H₂O: C: 52.63; H: 5.96; N: 15.34. Found: C: 52.62; H: 5.70; N: 15.32; major isomer (26.2). mp 225 - 230°C; Anal. Cald. for C₂₀H₂₆N₅O₅P + 0.5 H₂O: C: 52.63; H: 5.96; N: 15.34. Found: C: 52.74 ; H: 5.80; N: 15.32.

Following the above described procedures, N⁹-phenethyl-8-(2-(5-(5-hydroxyl-1,3 cyclohexyl)phosphono)furanyl)adenine (26.3) was also prepared. Anal. Cald. for C₂₃H₂₄N₅O₅P + 0.15 H₂O: C: 57.06 ; H: 5.06 ; N: 14.47. Found: C: 56.84; H: 4.83; N: 14.38.

Example 27.**Preparation of 3-(2-hydroxyethyl)phthalide.**

A solution of phthalide-3-acetic acid (1 mmol) in THF was treated with borane dimethylsulfide (1.5 mmol) at 0 °C for 1h, and 25 °C for 24 h. Extraction and chromatography gave 2-(3-phthalidyl)ethanol as a light yellow oil. TLC: R_f = 0.25, 50 % EtOAc - hexane.

Example 28.**Preparation of purine phosphonate amine salts**

A mixture of N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine (1 mmol) and tris(hydroxymethyl)aminomethane (1.05 mmol) in methanol is stirred at 25 °C for 24 h. Evaporation give N⁹-neopentyl-8-(2-(5-phosphono)furanyl)adenine tris(hydroxymethyl)aminomethane salt .

Examples of the methods of the present invention includes the following. It will be understood that these examples are exemplary and that the method of the invention is not limited solely to these examples.

For the purposes of clarity and brevity, chemical compounds in the following biological examples are referred to by synthetic example numbers.

Besides the following Examples, assays that may be useful for identifying compounds which inhibit gluconeogenesis include the following animal models of Diabetes:

- i. Animals with pancreatic b-cells destroyed by specific chemical cytotoxins such as Alloxan or Streptozotocin (e.g. the Streptozotocin-treated mouse, -rat, dog, and -monkey). Kodama, H., Fujita, M., Yamaguchi, I.,
10 *Japanese Journal of Pharmacology* 1994, 66, 331-336 (mouse); Youn, J.H., Kim, J.K., Buchanan, T.A., *Diabetes* 1994, 43, 564-571 (rat); Le Marchand, Y., Loten, E.G., Assimacopoulos-Jannet, F., et al., *Diabetes* 1978, 27, 1182-88 (dog); and Pitkin, R.M., Reynolds, W.A., *Diabetes* 1970, 19, 70-85 (monkey).
- ii. Mutant mice such as the C57BL/Ks db/db, C57BL/Ks ob/ob, and
15 C57BL/6J ob/ob strains from Jackson Laboratory, Bar Harbor, and others such as Yellow Obese, T-KK, and New Zealand Obese. Coleman, D.L., Hummel, K.P., *Diabetologia* 1967, 3, 238-248 (C57BL/Ks db/db); Coleman, D.L., *Diabetologia* 1978, 14, 141-148 (C57BL/6J ob/ob); Wolff, G.L., Pitot, H.C.,
20 *Genetics* 1973, 73, 109-123 (Yellow Obese); Dulin, W.E., Wyse, B.M., *Diabetologia* 1970, 6, 317-323 (T-KK); and Bielschowsky, M., Bielschowsky, F. *Proceedings of the University of Otago Medical School* 1953, 31, 29-31 (New Zealand Obese).
- iii. Mutant rats such as the Zucker fa/fa Rat rendered diabetic with Streptozotocin or Dexamethasone, the Zucker Diabetic Fatty Rat, and the Wistar
25 Kyoto Fatty Rat. Stolz, K.J., Martin, R.J. *Journal of Nutrition* 1982, 112, 997-1002 (Streptozotocin); Ogawa, A., Johnson, J.H., Ohnbede, M., McAllister, C.T., Irman, L., Alam, T., Unger, R.H., *The Journal of Clinical Investigation* 1992, 90, 497-504 (Dexamethasone); Clark, J.B., Palmer, C.J., Shaw, W.N., *Proceedings of the Society for Experimental Biology and Medicine* 1983, 173, 68-75 (Zucker
30 Diabetic Fatty Rat); and Idida, H., Shino, A., Matsuo, T., et al., *Diabetes* 1981, 30, 1045-1050 (Wistar Kyoto Fatty Rat).
- iv. Animals with spontaneous diabetes such as the Chinese Hamster, the Guinea Pig, the New Zealand White Rabbit, and non-human primates such as the Rhesus monkey and Squirrel monkey. Gerritsen, G.C., Connel, M.A.,
35 Blanks, M.C., *Proceedings of the Nutrition Society* 1981, 40, 237-245 (Chinese Hamster); Lang, C.M., Munger, B.L., *Diabetes* 1976, 25, 434-443 (Guinea Pig);

Conaway, H.H., Brown, C.J., Sanders, L.L. et al., *Journal of Heredity* **1980**, *71*, 179-186 (New Zealand White Rabbit); Hansen, B.C., Bodkin, M.L., *Diabetologia* **1986**, *29*, 713-719 (Rhesus monkey); and Davidson, I.W., Lang, C.M., Blackwell, W.L., *Diabetes* **1967**, *16*, 395-401 (Squirrel monkey).

- 5 v. Animals with nutritionally induced diabetes such as the Sand Rat, the Spiny Mouse, the Mongolian Gerbil, and the Cohen Sucrose-Induced Diabetic Rat. Schmidt-Nielsen, K., Hainess, H.B., Hackel, D.B., *Science* **1964**, *143*, 689-690 (Sand Rat); Gonet, A.E., Stauffacher, W., Pictet, R., et al., *Diabetologia* **1965**, *1*, 162-171 (Spiny Mouse); Boquist, L., *Diabetologia* **1972**, *8*, 274-282 (Mongolian Gerbil); and Cohen, A.M., Teitebaum, A., Salitemik, R., *Metabolism* **1972**, *21*, 235-240 (Cohen Sucrose-Induced Diabetic Rat).
- 10

- vi. Any other animal with one of the following or a combination of the following characteristics resulting from a genetic predisposition, genetic engineering, selective breeding, or chemical or nutritional induction: impaired
- 15 glucose tolerance, insulin resistance, hyperglycemia, obesity, accelerated gluconeogenesis, increased hepatic glucose output.

Example A: Inhibition of Human Liver FBPase

- E. coli* strain BL21 transformed with a human liver FBPase-encoding
- 20 plasmid was obtained from Dr. M. R. El-Maghrabi at the State University of New York at Stony Brook. hFBPase was typically purified from 10 liters of *E. coli* culture as described (M. Gidh-Jain et al., *The Journal of Biological Chemistry* **1994**, *269*, 27732-27738). Enzymatic activity was measured spectrophotometrically in reactions that coupled the formation of product
- 25 (fructose 6-phosphate) to the reduction of dimethylthiazoldiphenyltetrazolium bromide (MTT) via NADP and phenazine methosulfate (PMS) , using phosphoglucose isomerase and glucose 6-phosphate dehydrogenase as the coupling enzymes. Reaction mixtures (200 μ L) were made up in 96-well microtitre plates, and consisted of 50 mM Tris-HCl, pH 7.4, 100 mM KCl, 5 mM
- 30 EGTA, 2 mM $MgCl_2$, 0.2 mM NADP, 1 mg/mL BSA, 1 mM MTT, 0.6 mM PMS, 1 unit/mL phosphoglucose isomerase, 2 units/mL glucose 6-phosphate dehydrogenase, and 0.150 mM substrate (fructose 1,6-bisphosphate). Inhibitor concentrations were varied from 0.01 μ M to 10 μ M. Reactions were started by the addition of 0.002 units of pure hFBPase and were monitored for 7 minutes
- 35 at 590 nm in a Molecular Devices Plate Reader (37 $^{\circ}$ C).

The following Table depicts the IC₅₀ values for several compounds prepared in the Examples. AMP has an IC₅₀ value of 1.0 μ M in this assay.

Example	Compound		IC ₅₀ (human liver FBPase)
	Number		
5	2.1	5	μ M
	2.2	1.4	μ M
	2.3	3.3	μ M
10	2.7	0.8	μ M
	2.10	2	μ M
	2.13	4.5	μ M
	2.14	0.9	μ M
	2.16	1.4	μ M
15	4.1	100	μ M
	5.5	10	μ M
	8.1	23	μ M
	10.1	50	μ M
	11.1	0.7	μ M
20	12.1	13	μ M

Figure 2 shows the dose-dependent inhibition of hIFBPase by AMP, compound 2.7, compound 2.13, and compound 2.5.

***In vitro* inhibition of rat liver and mouse liver FBPase was also determined.**

E. coli strain BL21 transformed with a rat liver FBPase-encoding plasmid was obtained from Dr. M. R. El-Maghrabi at the State University of New York at Stony Brook, and purified as described (El-Maghrabi, M.R., and Pilgis, S.J. *Biochem. Biophys. Res. Commun.* 1991, 176, 137-144). Mouse liver FBPase was obtained by homogenizing freshly isolated mouse liver in 100 mM Tris-HCl buffer, pH 7.4, containing 1 mM EGTA, and 10 % glycerol. The homogenate was clarified by centrifugation, and the 45-75 % ammonium sulfate fraction prepared. This fraction was redissolved in the homogenization buffer and desalted on a PD-10 gel filtration column (Biorad) eluted with same. This partially purified fraction was used for enzyme assays. Both rat liver and mouse liver FBPase were assayed as described for human liver FBPase. Generally,

as reflected by the higher IC₅₀ values, the rat and mouse liver enzymes are less sensitive to inhibition by the compounds tested than the human liver enzyme.

The following Table depicts the IC₅₀ values for several compounds prepared in the Examples:

	Compound	IC ₅₀ Rat Liver	IC ₅₀ Mouse Liver
		μ M	μ M
10	2.1	100	>20
	2.2	>20	>20
	2.3	>20	>20
	2.7	1.25	55
	2.10	>20	>20
15	2.13	>20	>20
	2.14	>20	>20
	2.16	>20	>20
	4.1	>20	>20
	5.5	>20	>20
20	8.1	>20	>20
	10.1	>20	>20
	11.1	>20	>20
	12.1	20	>100

Example B: AMP Site Binding

To determine whether compounds bind to the allosteric AMP binding site of hIFBPase, the enzyme was incubated with radiolabeled AMP in the presence of a range of test compound concentrations. The reaction mixtures consisted of 25 mM ³H-AMP (54 mCi/mmol) and 0 -1000 mM test compound in 25 mM Tris-HCl, pH 7.4, 100 mM KCl and 1 mM MgCl₂. 1.45 mg of homogeneous FBPase (\pm 1 nmole) was added last. After a 1 minute incubation, AMP bound to FBPase was separated from unbound AMP by means of a centrifugal ultrafiltration unit ("Ultrafree-MC", Millipore) used according to the instructions of the manufacturer. The radioactivity in aliquots (100 μ l) of the upper compartment of the unit (the retentate, which contains enzyme and label) and the lower compartment (the filtrate, which contains unbound label) were quantified using a Beckman liquid scintillation counter. The amount of AMP bound to the enzyme was estimated by comparing the counts in the filtrate (the unbound label) to the total counts in the retentate.

As evident from Fig. 3, both aminoimidazolecarboxamideriboside monophosphate (ZMP) and compound 2.2 displaced AMP from hIFBPase in a

dose-dependent manner, indicating that they bind to the same site on the enzyme as AMP. As expected, compound 2.2, a more potent hFBPase inhibitor than ZMP (IC_{50} 's = 5 and 12 μ M, respectively), had a lower ED_{50} for AMP displacement than ZMP (35 vs 250 μ M).

5

Example C: AMP Site/Enzyme Selectivity

To determine the selectivity of compounds towards FBPase, effects of FBPase inhibitors on 5 key AMP binding enzymes were measured using the assays described below:

10

Adenosine Kinase: Human adenosine kinase was purified from an *E. coli* expression system as described by Spychala *et al.* (Spychala, J., Datta, N.S., Takabayashi, K., Datta, M., Fox, I.H., Gribbin, T., and Mitchell, B.S. *Proc. Natl. Acad. Sci. USA* 1996, 93, 1232-1237). Activity was measured essentially as described by Yamada *et al.* (Yamada, Y., Goto, H., Ogasawara, N. *Biochim. Biophys. Acta* 1988, 660, 36-43.) with a few minor modifications. Assay mixtures contained 50 mM TRIS-maleate buffer, pH 7.0, 0.1 % BSA, 1 mM ATP 1 mM $MgCl_2$, 1.0 μ M [$U-^{14}C$] adenosine (400-600 mCi/mmol) and varying duplicate concentrations of inhibitor. ^{14}C -AMP was separated from unreacted ^{14}C -adenosine by absorption to anion exchange paper (Whatman) and quantified by scintillation counting.

Adenosine Monophosphate Deaminase: Porcine heart AMPDA was purified essentially as described by Smiley *et al.* (Smiley, K.L., Jr, Berry, A.J., and Suelter, C.H. *J. Biol. Chem.* 1967, 242, 2502-2506) through the phosphocellulose step. Inhibition of AMPDA activity was determined at 37 °C in a 0.1 mL assay mixture containing inhibitor, ~0.005U AMPDA, 0.1 % bovine serum albumin, 10 mM ATP, 250 mM KCl, and 50 mM MOPS at pH 6.5. The concentration of the substrate AMP was varied from 0.125 - 10.0 mM. Catalysis was initiated by the addition of enzyme to the otherwise complete reaction mixture, and terminated after 5 minutes by injection into an HPLC system. Activities were determined from the amount of IMP formed during 5 minutes. IMP was separated from AMP by HPLC using a Beckman Ultrasil-SAX anion exchange column (4.6 mm x 25 cm) with an isocratic buffer system (12.5 mM potassium phosphate, 30 mM KCl, pH 3.5) and detected spectrophotometrically by absorbance at 254 nm.

35

- Phosphofructokinase:* Enzyme (rabbit liver) was purchased from Sigma. Activity was measured at 30 °C in reactions in which the formation of fructose 1,6-bisphosphate was coupled to the oxidation of NADH via the action of aldolase, triosephosphate isomerase, and α -glycerophosphate
- 5 dehydrogenase. Reaction mixtures (200 μ l) were made up in 96-well microtitre plates and were read at 340 nm in a Molecular Devices Microplate Reader. The mixtures consisted of 200 mM Tris-HCl pH 7.0, 2 mM DTT, 2 mM $MgCl_2$, 0.2 mM NADH, 0.2 mM ATP, 0.5 mM Fructose 6-phosphate, 1 unit aldolase/mL, 3 units/mL triosephosphate isomerase, and 4 units/mL α -glycerophosphate
- 10 dehydrogenase. Test compound concentrations ranged from 1 to 500 μ M. Reactions were started by the addition of 0.0025 units of phosphofructokinase and were monitored for 15 minutes.
- Glycogen Phosphorylase:* Enzyme (rabbit muscle) was purchased from Sigma.
- 15 Activity was measured at 37 °C in reactions in which the formation of glucose 1-phosphate was coupled to the reduction of NADP via phosphoglucomutase and glucose 6-phosphate dehydrogenase. Assays were performed on 96-well microtitre plates and were read at 340 nm on a Molecular Devices Microplate Reader. Reaction mixtures consisted of 20 mM imidazole, pH 7.4, 20 mM
- 20 $MgCl_2$, 150 mM potassium acetate, 5 mM potassium phosphate, 1 mM DTT, 1 mg/mL BSA, 0.1 mM NADP, 1 unit/mL phosphoglucomutase, 1 unit/mL glucose 6-phosphate dehydrogenase, 0.5 % glycogen. Test compound concentrations ranged from 1 to 500 μ M. Reactions were started by the addition of 17 μ g enzyme and were monitored for 20 minutes.
- 25
- Adenylate Kinase:* Enzyme (rabbit muscle) was purchased from Sigma. Activity was measured at 37 °C in reaction mixtures (100 μ l) containing 100 mM Hepes, pH 7.4, 45 mM $MgCl_2$, 1 mM EGTA, 100 mM KCl, 2 mg/mL BSA, 1 mM AMP and 2 mM ATP. Reactions were started by addition of 4.4 ng enzyme and
- 30 terminated after 5 minutes by addition of 17 μ l perchloric acid. Precipitated protein was removed by centrifugation and the supernatant neutralized by addition of 33 μ l 3 M KOH/3 M KH_2CO_3 . The neutralized solution was clarified by centrifugation and filtration and analyzed for ADP content (enzyme activity) by HPLC using a YMC ODS AQ column (25 X 4.6 cm). A gradient was run from
- 35 0.1 M KH_2PO_4 , pH 6, 8 mM tetrabutyl ammonium hydrogen sulfate to 75 % acetonitrile. Absorbance was monitored at 254 nM.

Compound 2.1, a 5 μ M hIFBPase inhibitor, was essentially inactive in all of the above described assays except for the AMP deaminase screen: half-maximal inhibition of AMP deaminase was observed at almost the same concentration as the IC_{50} for FBPase. Compound 2.7 (hIFBPase IC_{50} = 0.8 μ M), in addition to being essentially without effect on adenosine kinase, adenylate kinase, glycogen phosphorylase, and phosphofructokinase, was only a weak inhibitor of AMP deaminase (IC_{50} = 390 μ M). The data suggest that compound 2.7 binds to hIFBPase in a highly selective manner. The following Table gives the selectivity data for compounds 2.1 and 2.7.

SELECTIVITY

	<u>Compound 2.1</u>	<u>Compound 2.7</u>
FBPase (inh.)	5.0 μ M	0.8 μ M
15 Adenosine Kinase (inh.)	>>10	>>100
Adenylate Kinase (inh.)	>>500	>>500
AMP Deaminase (inh.)	6.7	390
Glycogen Phosphorylase (act.)	>>250	>>100
20 Phosphofructokinase (act.)	>>200	>>100

Example D: Inhibition of Gluconeogenesis in Rat Hepatocytes

Hepatocytes were prepared from overnight fasted Sprague-Dawley rats (250-300 g) according to the procedure of Berry and Friend (Berry, M.N., Friend, D.S., *J. Cell. Biol.* **1969**, *43*, 506-520) as modified by Groen (Groen, A.K., Sips, H.J., Vervoorn, R.C., Tager, J.M., *Eur. J. Biochem.* **1982**, *122*, 87-93). Hepatocytes (75 mg wet weight/mL) were incubated in 1 mL Krebs-bicarbonate buffer containing 10 mM Lactate, 1 mM pyruvate, 1 mg/mL BSA, and test compound concentrations from 1 to 500 μ M. Incubations were carried out in a 95 % oxygen, 5 % carbon dioxide atmosphere in closed, 50-mL Falcon tubes submerged in a rapidly shaking water bath (37 °C). After 1 hour, an aliquot (0.25 mL) was removed, transferred to an Eppendorf tube and centrifuged. 50 μ L of supernatant was then assayed for glucose content using a Sigma Glucose Oxidase kit as per the manufacturer's instructions.

Compound 2.1 and compound 2.7 inhibited glucose production from lactate/pyruvate in isolated rat hepatocytes in a dose-dependent manner, with IC_{50} 's of 90 and 4.5 μ M, respectively (Figure 5). IC_{50} 's for other select compounds in this assay are shown in the Table below:

5		<u>Compound</u>	<u>IC_{50} Glucose Production, μM</u>
		2.2	90
		2.6	18
		2.10	24
10		2.13	50
		2.14	7.5
		2.16	12
		16.4	3

15 FBPase from rat liver is less sensitive to AMP than that from human liver. IC_{50} values are correspondingly higher in rat hepatocytes than would be expected in human hepatocytes.

20 **Example E: Effect of Compound 2.7 on gluconeogenesis from Dihydroxyacetone in Rat Hepatocytes: Glucose Production Inhibition and Fructose 1,6-bisphosphate Accumulation**

Isolated rat hepatocytes were prepared as described in Example D and incubated under the identical conditions described except that lactate/pyruvate was replaced by 10 mM dihydroxyacetone, a substrate which feeds into the gluconeogenic pathway at a step just prior to FBPase. Reactions were
25 terminated by removing an aliquot (250 μ L) of cell suspension and spinning it through a layer of oil (0.8 mL silicone/mineral oil, 4/1) into a 10 % perchloric acid layer (100 μ L). After removal of the oil layer, the acidic cell extract layer was neutralized by addition of 1/3rd volume of 3 M KOH/3 M KH_2CO_3 . After
30 thorough mixing and centrifugation, the supernatant was analyzed for glucose content as described in Example D, and also for fructose-1,6-bisphosphate. Fructose-1,6-bisphosphate was assayed spectrophotometrically by coupling its enzymatic conversion to glycerol 3-phosphate to the oxidation of NADH, which was monitored at 340 nm. Reaction mixtures (1 mL) consisted of 200 mM Tris-HCl, pH 7.4, 0.3 mM NADH, 2 units/mL glycerol 3-phosphate dehydrogenase, 2
35 units/mL triosephosphate isomerase, and 50-100 μ L cell extract. After a 30

minute preincubation at 37 °C, 1 unit/mL of aldolase was added and the change in absorbance measured until a stable value was obtained. 2 moles of NADH are oxidized in this reaction per mole of fructose-1,6-bisphosphate present in the cell extract.

- 5 As shown in Figure 4A, compound 2.7 inhibited glucose production from dihydroxyacetone in rat hepatocytes (IC_{50} approx. 5 μ M) as effectively as from lactate pyruvate (IC_{50} 4.5 μ M, Figure 5). This data confirms that the site of action of the compound is in the last four steps of the gluconeogenic pathway. The dose-dependent accumulation of fructose-1,6-bisphosphate (the substrate of
- 10 FBPase) that occurs upon cell exposure to compound 2.7 (Figure 4B) is consistent with the inhibition of FBPase, the second to last enzyme in the pathway.

Example F: Blood Glucose Lowering in Fasted Rats

- 15 Sprague Dawley rats (250-300 g) were fasted for 18 hours and then dosed intraperitoneally either with saline or with 35, 45, and 60 mg/kg compound 16.4, a prodrug of compound 2.7. The vehicle used for drug administration was dimethylsulfoxide. Blood samples were obtained from the tail vein of conscious animals just prior to injection and then at half-hourly
- 20 intervals. Blood glucose was measured using a HemoCue Inc. glucose analyzer according to the instructions of the manufacturer.

Figure 6 shows the profound glucose lowering elicited by treatment with compound 16.4. The duration of action was dose-dependent and ranged from 2 to 6 hours.

25

Example G: Analysis of Drug Levels and Liver Fructose-1,6-bisphosphate Accumulation In Rats

- Sprague-Dawley rats (250-300 g) were fasted for 18 hours and then dosed intraperitoneally either with saline ($n = 3$) or 20 mgs/kg compound 2.7 (n
- 30 $= 4$). The vehicle used for drug administration was 10 mM bicarbonate. One hour post injection rats were anesthetized with halothane and a liver biopsy (approx. 1 g) was taken as well as a blood sample (2 mL) from the posterior vena cava. A heparin flushed syringe and needle were used for blood collection. The liver sample was immediately homogenized in ice-cold 10% perchloric acid (3 mL), centrifuged, and the supernatant neutralized with 1/3rd
- 35 volume of 3 M KOH/3 M KH_2CO_3 . Following centrifugation and filtration, 50 μ L

of the neutralized extract was analyzed for compound 2.7 content by HPLC. A reverse phase YMC ODS AQ column (250 X 4.6 cm) was used and eluted with a gradient from 10 mM sodium phosphate pH 5.5 to 75 % acetonitrile.

Absorbance was monitored at 310 nm. The concentration of fructose-1,6-

5 bisphosphate in liver was also quantified using the method described in Example E. Blood glucose was measured in the blood sample as described in Example F. Plasma was then quickly prepared by centrifugation and extracted by addition of methanol to 60 % (v/v). The methanolic extract was clarified by centrifugation and filtration and then analyzed by HPLC as described above.

10 Compound 2.7 lowered blood glucose from 82 ± 3 to 28 ± 9.9 mg/dL within one hour (Figure 7). Drug levels measured in plasma and liver were 38.5 ± 7 μ M and 51.3 ± 10 nmoles/ g, respectively. As shown in Figure 8, a 10-fold elevation of fructose-1,6-bisphosphate levels was found in the livers from the drug-treated group, consistent with the inhibition of glucose production at the
15 level of FBPase in the gluconeogenic pathway.

Example H: Blood Glucose Lowering in Zucker Diabetic Fatty Rats

Zucker Diabetic Fatty rats purchased at 7 weeks of age were used at age 16 weeks in the 24-hour fasted state. The rats were purchased from Genetics
20 Models Inc. and fed the recommended Purina 5008 diet (6.5 % fat). Their fasting hyperglycemia at 24 hours ranged from 150 mg/dL to 310 mg/dL blood glucose.

Compound 2.7 was administered at a dose of 50 mg/kg by intraperitoneal injection (n = 6). The stock solution was made up at 25 mg/mL in
25 deionized water and adjusted to neutrality by dropwise addition of 5 N NaOH. 5 control animals were dosed with saline. Blood glucose was measured at the time of dosing and 2 hours post dose as described in Example F.

As shown in Figures 9A and 9B, blood glucose was lowered in the drug-treated group by an average of almost 20 % ($p < 0.0001$ relative to the control
30 animals).

Example I: Inhibition of Gluconeogenesis in Zucker Diabetic Fatty Rats

Three 20-week old Zucker Diabetic Fatty rats were dosed with compound
35 2.7 and three with saline as described in Example H. Fifteen minutes post-injection, the animals were anesthetized with sodium pentobarbitol (30 mgs,

i.p.) and ^{14}C -bicarbonate (20 μCi /100 g of body weight) was administered via the tail vein. Blood samples (0.6 mL) were obtained by cardiac puncture 10 and 20 minutes post tracer injection. Blood (0.5 mL) was diluted into 6 mL deionized water and protein precipitated by addition of 1 mL zinc sulfate (0.3 N) and 1 mL barium hydroxide (0.3 N). The mixture was centrifuged (20 minutes, 1000X g) and 5 mL of the resulting supernatant was then combined with 1 g of a mixed bed ion exchange resin (1 part AG 50W-X8, 100-200 mesh, hydrogen form and 2 parts of AG 1-X8, 100-200 mesh, acetate form) to separate ^{14}C -bicarbonate from ^{14}C -glucose. The slurry was shaken at room temperature for four hours and then allowed to settle. An aliquot of the supernatant (0.5 mL) was then counted in 5 mL scintillation cocktail.

As shown in Figure 10, compound 2.7 reduced the incorporation of ^{14}C -bicarbonate into glucose by 75 %; therefore gluconeogenesis was clearly inhibited by the drug.

15

Example J: Blood Glucose Lowering in Streptozotocin-treated Rats

Diabetes is induced in male Sprague-Dawley rats (250-300g) by intraperitoneal injection of 55 mg/kg streptozotocin (Sigma Chemical Co.). Six days later, 24 animals are selected with fed blood glucose values (8 am) between 350 and 600 mg/dL and divided into two statistically equivalent groups. Blood glucose is measured in blood obtained from a tail vein nick by means of a HemoCue Inc. (Mission Viejo, CA) glucose analyzer. One group of 12 will subsequently receive inhibitor (100 mg/kg intraperitoneally) and the other 12 ("controls") an equivalent volume of saline. Food is removed from the animals. Blood glucose is measured in each animal four hours after dosing, and a second dose of drug or saline is then administered. Four hours later, a final blood glucose measurement is made.

Example K: Evaluation of Compound 16.4 as a Prodrug in Rat Hepatocytes - Intracellular Delivery of Compound 2.7

Rat hepatocytes were prepared and incubated as in Example D, except that the test compound, 16.4, was added to yield a final concentration of 10 μM . Aliquots of the cell suspension were taken at 0, 5, 10, 20, 30, 45, and 60 minutes after drug exposure. The cells were extracted and analyzed for compound 2.7 content by HPLC as described in Example L. Absorbance of the HPLC column eluate was monitored at 310 nm. Quantitation of intracellular

compound 2.7 was done by comparison to authentic standards of known concentration. As shown in Figure 11A, compound 16.4 rapidly delivered high levels of compound 2.7 into the hepatocytes; a concentration of approximately 80 nmoles/g was achieved within 10 minutes. These data indicate that

5 compound 16.4 readily penetrates cells and is efficiently de-esterified to the parent compound, 2.7, intracellularly. Furthermore, as shown in Figure 11B, compound 16.4 inhibited glucose production in rat hepatocytes.

10 **Example L: Estimation of the oral bioavailability of prodrugs of phosphonic acids:**

Prodrugs were dissolved in 10 % ethanol/90 % polyethylene glycol (mw 400) and administered by oral gavage at doses of approximately 20 or 40 mg/kg parent compound equivalents to 6-hour fasted, Sprague Dawley rats (220-240

15 g). The rats were subsequently placed in metabolic cages and urine was collected for 24 hours. The quantity of parent compound excreted into urine was determined by HPLC analysis. An ODS column eluted with a gradient from potassium phosphate buffer, pH 5.5 to acetonitrile was employed for these measurements. Detection was at 310-325 nm. The percentage oral

20 bioavailability was estimated by comparison of the recovery in urine of the parent compound generated from the prodrug, to that recovered in urine 24 hours after intravenous administration of unsubstituted parent compound at approximately 10 mg/kg. Parent compounds were typically dissolved in dimethyl sulfoxide, and administered via the tail vein in animals that were briefly

25 anesthetized with halothane.

For compound 16.4, a prodrug of compound 2.7, 6.2 % of an oral dose of approximately 20 mg/kg was recovered in urine. For compound 2.7, 76.8 % of an intravenous dose of approximately 10 mg/kg was recovered. The oral bioavailability of compound 16.4 was therefore calculated to be 6.2/76.8, or


30 approximately 8 %. The oral bioavailability of compound 16.5 was also estimated following the above described protocol to be 5.3 %.

**Example 1. M. Glucose low ring following oral administration of
FBPase Inhibitors:**

FBPase inhibitor was administered by oral gavage at doses of 30, 100 and 250
5 mg/kg to 18-hour fasted, Sprague Dawley rats (250-300g; n= 4-5/group). The
compound was prepared in deionized water, adjusted to neutrality with sodium
hydroxide, and brought into solution by sonication prior to administration. Blood
glucose was measured immediately prior to dosing, and at 1 hour intervals
thereafter. Blood samples were obtained from the tail vein, and measurements
10 made by means of a Hemocue glucose analyzer (Hemocue Inc, Mission Viejo,
California) used according to the manufacturer's instructions.

We claim:

1. A compound of formula 1:



5

wherein

- A is selected from the group consisting of $-NR^8_2$, $NHSO_2R^3$, $-OR^5$, $-SR^5$, halogen, lower alkyl, $-CON(R^4)_2$, guanidine, amidine, $-H$, and perhaloalkyl;


- E is selected from the group consisting of $-H$, halogen, lower alkylthio, lower perhaloalkyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, $-CN$, and $-NR^7_2$;

- X is selected from the group consisting of alkylamino, alkyl, alkenyl, alkynyl, alkyl(carboxyl), alkyl(hydroxy), alkyl(phosphonate), alkyl(sulfonate), aryl, alkylaminoalkyl, alkoxyalkyl, alkylthioalkyl, alkylthio, alicyclic, 1,1-dihaloalkyl, carbonylalkyl, aminocarbonylamino, alkylaminocarbonyl, alkylcarbonylamino, aralkyl, and alkylaryl, all optionally substituted; or together with Y forms a cyclic group including cyclic alkyl, heterocyclic, and aryl;

- Y is selected from the group consisting of $-H$, alkyl, alkenyl, alkynyl, aryl, alicyclic, aralkyl, aryloxyalkyl, alkoxyalkyl, $-C(O)R^3$, $-S(O)_2R^3$, $-C(O)-OR^3$, $-CONHR^3$, $-NR^2_2$, and $-OR^3$, all except H are optionally substituted; or together with X forms a cyclic group including aryl, cyclic alkyl, and heterocyclic;

- R^1 is independently selected from the group consisting of $-H$, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or thiocarbonate, $-C(R^2)_2$ -aryl, alkylaryl, $-C(R^2)_2OC(O)NR^2_2$, $-NR^2-C(O)-R^3$, $-C(R^2)_2-OC(O)R^3$, $C(R^2)_2-O-C(O)OR^3$, $-C(R^2)_2OC(O)SR^3$, alkyl-S-C(O) R^3 , alkyl-S-S-alkylhydroxy, and alkyl-S-S-S-alkylhydroxy, or together R^1 and R^1 are -alkyl-S-S-alkyl to form a cyclic group, or together R^1 and R^1 are

30



wherein

V and W are independently selected from the group consisting of
 5 hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or

together V and Z are connected to form a cyclic group containing 3-5
 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy,
 alkoxy, alkoxy, or aryloxy, or aryloxy attached to a carbon atom that is three atoms
 10 from an oxygen attached to the phosphorus; or

together V and W are connected to form a cyclic group containing 3
 carbon atoms substituted with hydroxy, acyloxy, alkoxy, alkoxy,
 alkylthio, hydroxymethyl, and aryloxy attached to a carbon atom
 that is three atoms from an oxygen attached to the phosphorus;

15 Z is selected from the group consisting of $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOR}^3$,
 $-\text{CH}_2\text{OC(O)SR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{SR}^3$, $-\text{S(O)R}^3$, $-\text{CH}_2\text{N}_3$, $-\text{CH}_2\text{NR}^2_2$, $-\text{CH}_2\text{Ar}$,
 $-\text{CH(Ar)OH}$, $-\text{CH}(\text{CH}=\text{CR}^2\text{R}^2)\text{OH}$, $-\text{CH}(\text{C}\equiv\text{CR}^2)\text{OH}$, and $-\text{R}^2$;

with the provisos that:

- a) V, Z, W are not all $-\text{H}$; and
 20 b) when Z is $-\text{R}^2$, then at least one of V and W is not $-\text{H}$ or $-\text{R}^9$;
 R^2 is selected from the group consisting of R^3 and $-\text{H}$;
 R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and
 aralkyl;

R^4 is independently selected from the group consisting of $-\text{H}$, lower alkyl,
 25 lower alicyclic, lower aralkyl, and lower aryl;

R^5 is selected from the group consisting of lower alkyl, lower aryl, lower
 aralkyl, and lower alicyclic;

R^6 is independently selected from the group consisting of $-\text{H}$, and lower
 alkyl;

30 R^7 is independently selected from the group consisting of $-\text{H}$, lower alkyl,
 lower alicyclic, lower aralkyl, lower aryl, and $-\text{C(O)R}^{10}$;

R^8 is independently selected from the group consisting of -H, lower alkyl, lower aralkyl, lower aryl, lower alicyclic, $-C(O)R^{10}$, or together they form a bidentate alkyl;

5 R^9 is selected from the group consisting of alkyl, aralkyl, and alicyclic;

R^{10} is selected from the group consisting of -H, lower alkyl, $-NH_2$, lower aryl, and lower perhaloalkyl;

10 R^{11} is selected from the group consisting of alkyl, aryl, -OH, $-NH_2$ and $-OR^3$; and pharmaceutically acceptable prodrugs and salts thereof.

2. The compounds of claim 1 with the proviso that R^1 is not lower alkyl of 1-4 carbon atoms.

15 3. The compounds of claim 1 wherein A is selected from the group consisting of $-NR^8_2$, halogen, lower alkyl, lower perhaloalkyl, and lower alkoxy.

20 4. The compounds of claim 1 wherein E is -H, halogen, lower perhaloalkyl, -CN, lower alkyl, lower alkoxy, and lower alkylthio.

25 5. The compounds of claim 1 wherein X is selected from the group consisting of alkylamino, alkyl, alkynyl, alkoxyalkyl, alkylthio, aryl, 1,1-dihaloalkyl, carbonylalkyl, heteroaryl, alkylcarbonylamino, and alkylaminocarbonyl.


6. The compounds of claim 5 wherein X is alkyl substituted with 1 to 3 substituents selected from the group consisting of halogen, phosphonate, $-CO_2H$, $-SO_3H$, and -OH.

30 7. The compounds of claim 5 wherein X is selected from the group consisting of alkylaminocarbonyl, alkoxyalkyl, and heteroaryl.

8. The compounds of claim 7 wherein X is selected from the group consisting of methoxymethyl and optionally substituted furanyl.

9. The compounds of claim 1 wherein Y is selected from the group consisting of aralkyl, aryl, alicyclic, and alkyl.

10. The compounds of claim 1 wherein each R^1 is independently
 5 selected from the group consisting of -H, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or thiocarbonate, optionally substituted phenyl, optionally substituted benzyl, optionally substituted alkylaryl, $-C(R^2)_2OC(O)R^3$, $C(R^2)_2-O-C(O)OR^3$, $-C(R^2)_2-OC(O)SR^3$, -alkyl-S-C(O) R^3 , alkyl-S-S-alkylhydroxyl, and -alkyl-S-S-S-alkylhydroxy, or together R^1 and R^1 are alkyl-S-S-alkyl to form
 10 a cyclic group, or R^1 and R^1 together are



15 wherein

V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^3$; or


- together V and Z are connected to form a cyclic group containing 3-5
 20 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxycarboxy, or aryloxycarboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

- together V and W are connected to form a cyclic group containing 3
 carbon atoms substituted with hydroxy, acyloxy, alkoxycarboxy,
 25 alkylthiocarboxy, hydroxymethyl, and aryloxycarboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;

Z is selected from the group consisting of $-CH_2OH$, $-CH_2OCOR^3$, $-CH_2OC(O)SR^3$, $-CH_2OCO_2R^3$, $-SR^3$, $-S(O)R^3$, $-CH_2N_3$, $-CH_2NR^2_2$, $-CH_2Ar$, $-CH(Ar)OH$, $-CH(CH=CR^2R^2)OH$, $-CH(C\equiv CR^2)OH$, and $-R^2$;

with the provisos that:

- a) V, Z, W are not all -H; and
 - b) when Z is -R², then at least one of V and W is not -H or -R⁹;
R² is selected from the group consisting of R³ and -H;
- 5 R³ is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl; and
R⁹ is selected from the group consisting of alkyl, aralkyl, and alicyclic.
- 10 11. The compounds of claim 10 wherein each R¹ is independently selected from the group consisting of optionally substituted phenyl, optionally substituted benzyl, -C(R²)₂OC(O)R³, -C(R²)₂OC(O)OR³, and -H.
12. The compounds of claim 10 wherein R¹ is H.
- 15 13. The compounds of claim 10 wherein at least one R¹ is aryl, or -C(R²)₂-aryl.
14. The compounds of claim 10 wherein at least one R¹ is -C(R²)₂-
20 OC(O)R³, -C(R²)₂-OC(O)OR³, -C(R²)₂-OC(O)SR³.
15. The compounds of claim 10 wherein at least one R¹ is alkyl-S-S-alkylhydroxyl, -alkyl-S-C(O)R³, and -alkyl-S-S-S-alkylhydroxy, or together R¹ and R¹ are alkyl-S-S-alkyl to form a cyclic group.
- 25 16. The compounds of claim 10 wherein together R¹ and R¹ are



- 30 to form a cyclic group;
wherein

V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or

5 together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxy, alkoxy, or aryloxy, attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

10 together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxy, alkoxy, alkylthiocarboxy, hydroxymethyl, and aryloxy, attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;

Z is selected from the group consisting of $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOR}^3$, $-\text{CH}_2\text{OC(O)SR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{SR}^3$, $-\text{S(O)R}^3$, $-\text{CH}_2\text{N}_3$, $-\text{CH}_2\text{NR}^2_2$, $-\text{CH}_2\text{Ar}$, $-\text{CH(Ar)OH}$, $-\text{CH}(\text{CH}=\text{CR}^2\text{R}^2)\text{OH}$, $-\text{CH}(\text{C}\equiv\text{CR}^2)\text{OH}$, and $-\text{R}^2$;

15 with the provisos that:

a) V, Z, W are not all $-\text{H}$; and

b) when Z is $-\text{R}^2$, then at least one of V and W is not $-\text{H}$ or $-\text{R}^9$;

R^2 is selected from the group consisting of R^3 and $-\text{H}$;

20 R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl; and

R^9 is selected from the group consisting of alkyl, aralkyl, and alicyclic.

25 17. The compounds of claim 16 wherein V and W both form a 6-membered carbocyclic ring substituted with 0-4 groups, selected from the group consisting of hydroxy, acyloxy, alkoxy, alkoxy, and alkoxy; and Z is R^2 .

30 18. The compounds of claim 16 wherein V and W are hydrogen; and Z is selected from the group consisting of hydroxyalkyl, acyloxyalkyl, alkoxyalkyl, and alkoxyalkyl.

35 19. The compounds of claim 16 wherein V and W are independently selected from the group consisting of hydrogen, optionally substituted aryl, and optionally substituted heteroaryl, with the proviso that at least one of V and W is optionally substituted aryl or optionally substituted heteroaryl.

20. The compounds of claim 1 wherein together R¹ and R¹ are optionally substituted lactones attached at the omega position.

21. The compounds of claim 10 wherein R¹ is alicyclic where the cyclic moiety contains carbonate or thiocarbonate.

22. The compounds of claim 21 wherein together R¹ and R¹ are optionally substituted 2-oxo-1,3-dioxolenes attached through a methylene to the phosphorus oxygen.

10


23. The compounds of claim 1 wherein A is selected from the group consisting of -NR⁸₂, and halogen;

E is selected from the group consisting of -H, halogen, -CN, lower alkyl, lower perhaloalkyl, lower alkoxy, and lower alkylthio;

15 X is selected from the group consisting of alkylamino, alkyl, alkoxyalkyl, alkynyl, alkylthio, aryl, heteroaryl, alkylcarbonylamino, 1,1-dihaloalkyl, carbonylalkyl, alkyl(OH), alkyl(sulfonate), and alkylaminocarbonyl; and R⁴ and R⁷ are selected from the group consisting of -H, and lower alkyl.

20 24. The compounds of claim 23 wherein Y is selected from the group consisting of aralkyl, aryl, alicyclic, and alkyl.

25. The compound of claim 24 wherein R¹ and R¹ together are



25

V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and -R⁹; or

together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxycarboxy, or aryloxy-carboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

- 5 together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxycarboxy, alkylthiocarboxy, hydroxymethyl, and aryloxy-carboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;

10 Z is selected from the group consisting of $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOR}^3$, $-\text{CH}_2\text{OC(O)SR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{SR}^3$, $-\text{S(O)R}^3$, $-\text{CH}_2\text{N}_3$, $-\text{CH}_2\text{NR}^2_2$, $-\text{CH}_2\text{Ar}$, $-\text{CH(Ar)OH}$, $-\text{CH}(\text{CH}=\text{CR}^2\text{R}^2)\text{OH}$, $-\text{CH}(\text{C}\equiv\text{CR}^2)\text{OH}$, and $-\text{R}^2$;

with the provisos that:

- a) V, Z, W are not all -H; and
 b) when Z is $-\text{R}^2$, then at least one of V and W is not -H or $-\text{R}^9$;
 15 R^2 is selected from the group consisting of R^3 and -H;
 R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl; and
 R^9 is selected from the group consisting of alkyl, aralkyl, and alicyclic.

20

26. The compounds of claim 23 wherein A is $-\text{NR}^8_2$;

E is selected from the group consisting of -H, -Cl, and $-\text{SCH}_3$; and X is selected from the group consisting of optionally substituted furanyl and alkoxyalkyl.

25

27. The compounds of claim 26 wherein A is $-\text{NH}_2$; E is selected from the group consisting of -H, -Cl, and $-\text{SCH}_3$;

X is selected from the group consisting of 2,5-furanyl, and methoxymethyl; and

30

Y is lower alkyl.

28. The compound of claim 27 wherein E is -H, X is 2,5-furanyl, and Y is neopentyl.

35

29. The compound of claim 28 wherein R^1 is $-\text{CH}_2\text{O}-\text{C(O)}-\text{C}(\text{CH}_3)_3$ or its HCl salt.

30. The compound of claim 27 wherein E is -SCH₃, X is 2,5-furanyl, and Y is isobutyl.


5 31. The compound of claim 30 wherein R¹ is -CH₂O-C(O)-C(CH₃)₃ or its HCl salt.

32. The compound of claim 27 wherein E is -H, X is 2,5-furanyl, and Y is 1-(3-chloro-2,2-dimethyl)-propyl.

10

33. The compound of claim 32 wherein R¹ is -CH₂O-C(O)-C(CH₃)₃ or its HCl salt.

34. A method of treating an animal for diabetes mellitus, comprising
15 administering to said animal a therapeutically effective amount of a compound of formula (1):



20

wherein

A is selected from the group consisting of -NR⁸₂, NHSO₂R³, -OR⁵, -SR⁵, halogen, lower alkyl, -CON(R⁴)₂, guanidine, amidine, -H, and perhaloalkyl;

25 E is selected from the group consisting of -H, halogen, lower alkylthio, lower perhaloalkyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, -CN, and -NR⁷₂;

X is selected from the group consisting of alkylamino, alkyl, alkenyl, alkynyl, alkyl(carboxyl), alkyl(hydroxy), alkyl(phosphonate), alkyl(sulfonate),
30 aryl, alkylaminoalkyl, alkoxyalkyl, alkylthioalkyl, alkylthio, alicyclic, 1,1-dihaloalkyl, carbonylalkyl, aminocarbonylamino, alkylaminocarbonyl,

alkylcarbonylamino, aralkyl, and alkylaryl, all optionally substituted; or together with Y forms a cyclic group including cyclic alkyl, heterocyclic, and aryl;


Y is selected from the group consisting of -H, alkyl, alkenyl, alkynyl, aryl, alicyclic, aralkyl, aryloxyalkyl, alkoxyalkyl, $-C(O)R^3$, $-S(O)_2R^3$, $-C(O)-OR^3$,

- 5 $-CONHR^3$, $-NR^2_2$, and $-OR^3$, all except H are optionally substituted; or together with X forms a cyclic group including aryl, cyclic alkyl, and heterocyclic;

R^1 is independently selected from the group consisting of

-H, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or thiocarbonate, $-C(R^2)_2$ -aryl, alkylaryl, $-C(R^2)_2OC(O)NR^2_2$,

- 10 $-NR^2-C(O)-R^3$, $-C(R^2)_2-OC(O)R^3$, $C(R^2)_2-O-C(O)OR^3$, $-C(R^2)_2OC(O)SR^3$, alkyl-S- $C(O)R^3$, alkyl-S-S-alkylhydroxy, and alkyl-S-S-S-alkylhydroxy, or together R^1 and R^1 are -alkyl-S-S-alkyl to form a cyclic group, or together R^1 and R^1 are



15

wherein

V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or

20

together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxy, alkoxy, or aryloxy, or aryloxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

25

together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxy, alkoxy, alkylthiocarboxy, hydroxymethyl, and aryloxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;

Z is selected from the group consisting of $-CH_2OH$, $-CH_2OCOR^3$,

30

$-CH_2OC(O)SR^3$, $-CH_2OCO_2R^3$, $-SR^3$, $-S(O)R^3$, $-CH_2N_3$, $-CH_2NR^2_2$, $-CH_2Ar$, $-CH(Ar)OH$, $-CH(CH=CR^2R^2)OH$, $-CH(C\equiv CR^2)OH$, and $-R^2$;

with the provisos that:

a) V, Z, W are not all -H; and

b) when Z is -R², then at least one of V and W is not -H or -R⁹;

R² is selected from the group consisting of R³ and -H;

5 R³ is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl;

R⁴ is independently selected from the group consisting of -H, lower alkyl, lower alicyclic, lower aralkyl, and lower aryl;

10 R⁵ is selected from the group consisting of lower alkyl, lower aryl, lower aralkyl, and lower alicyclic;

R⁶ is independently selected from the group consisting of -H, and lower alkyl;

R⁷ is independently selected from the group consisting of -H, lower alkyl, lower alicyclic, lower aralkyl, lower aryl, and -C(O)R¹⁰;

15 R⁸ is independently selected from the group consisting of -H, lower alkyl, lower aralkyl, lower aryl, lower alicyclic, -C(O)R¹⁰, or together they form a bidendate alkyl;

R⁹ is selected from the group consisting of alkyl, aralkyl, and alicyclic;


20 R¹⁰ is selected from the group consisting of -H, lower alkyl, -NH₂, lower aryl, and lower perhaloalkyl;

R¹¹ is selected from the group consisting of alkyl, aryl, -OH, -NH₂ and -OR³; and

pharmaceutically acceptable prodrugs and salts thereof.

25

35. A method of lowering blood glucose levels in an animal in need thereof, comprising administering to said animal a pharmaceutically acceptable amount of a compound of formula (1):



30

wherein


A is selected from the group consisting of $-NR^8$, $NHSO_2R^3$, $-OR^5$, $-SR^5$, halogen, lower alkyl, $-CON(R^4)_2$, guanidine, amidine, $-H$, and perhaloalkyl;

5 E is selected from the group consisting of $-H$, halogen, lower alkylthio, lower perhaloalkyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, $-CN$, and $-NR^7$;

X is selected from the group consisting of alkylamino, alkyl, alkenyl, alkynyl, alkyl(carboxyl), alkyl(hydroxy), alkyl(phosphonate), alkyl(sulfonate),
10 aryl, alkylaminoalkyl, alkoxyalkyl, alkylthioalkyl, alkylthio, alicyclic, 1,1-dihaloalkyl, carbonylalkyl, aminocarbonylamino, alkylaminocarbonyl, alkylcarbonylamino, aralkyl, and alkylaryl, all optionally substituted; or together with Y forms a cyclic group including cyclic alkyl, heterocyclic, and aryl;

Y is selected from the group consisting of $-H$, alkyl, alkenyl, alkynyl, aryl,
15 alicyclic, aralkyl, aryloxyalkyl, alkoxyalkyl, $-C(O)R^3$, $-S(O)_2R^3$, $-C(O)-OR^3$, $-CONHR^3$, $-NR^2$, and $-OR^3$, all except H are optionally substituted; or together with X forms a cyclic group including aryl, cyclic alkyl, and heterocyclic;

R^1 is independently selected from the group consisting of $-H$, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or
20 thiocarbonate, $-C(R^2)_2$ -aryl, alkylaryl, $-C(R^2)_2OC(O)NR^2$, $-NR^2-C(O)-R^3$, $-C(R^2)_2-OC(O)R^3$, $C(R^2)_2-O-C(O)OR^3$, $-C(R^2)_2OC(O)SR^3$, alkyl-S-C(O) R^3 , alkyl-S-S-alkylhydroxy, and alkyl-S-S-S-alkylhydroxy, or together R^1 and R^1 are $-alkyl-S-S-alkyl$ to form a cyclic group, or together R^1 and R^1 are



25

wherein

V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or
30

together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxycarboxy, or aryloxy-carboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

- 5 together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxycarboxy, alkylthiocarboxy, hydroxymethyl, and aryloxy-carboxy attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;

Z is selected from the group consisting of $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOR}^3$,
10 $-\text{CH}_2\text{OC(O)SR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{SR}^3$, $-\text{S(O)R}^3$, $-\text{CH}_2\text{N}_3$, $-\text{CH}_2\text{NR}^2_2$, $-\text{CH}_2\text{Ar}$,
 $-\text{CH(Ar)OH}$, $-\text{CH}(\text{CH}=\text{CR}^2\text{R}^2)\text{OH}$, $-\text{CH}(\text{C}\equiv\text{CR}^2)\text{OH}$, and $-\text{R}^2$;

with the provisos that:

- a) V, Z, W are not all -H; and
b) when Z is $-\text{R}^2$, then at least one of V and W is not -H or $-\text{R}^8$;
15 R^2 is selected from the group consisting of R^3 and -H;
 R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl;

R^4 is independently selected from the group consisting of -H, lower alkyl, lower alicyclic, lower aralkyl, and lower aryl;

- 20 R^5 is selected from the group consisting of lower alkyl, lower aryl, lower aralkyl, and lower alicyclic;

R^6 is independently selected from the group consisting of -H, and lower alkyl;

- R^7 is independently selected from the group consisting of -H, lower alkyl,
25 lower alicyclic, lower aralkyl, lower aryl, and $-\text{C(O)R}^{10}$;

R^8 is independently selected from the group consisting of -H, lower alkyl, lower aralkyl, lower aryl, lower alicyclic, $-\text{C(O)R}^{10}$, or together they form a bidendate alkyl;

- R^9 is selected from the group consisting of alkyl, aralkyl, and
30 alicyclic;


R^{10} is selected from the group consisting of -H, lower alkyl, -NH₂, lower aryl, and lower perhaloalkyl;

R^{11} is selected from the group consisting of alkyl, aryl, -OH, -NH₂ and -OR³; and

5 pharmaceutically acceptable prodrugs and salts thereof.

36. A method of inhibiting FB Pase at the AMP site in patients in need thereof, comprising administering to said patients an FB Pase inhibitory amount of a compound of formula (1):

10



wherein

A is selected from the group consisting of -NR⁸₂, NHSO₂R³,
15 -OR⁵, -SR⁵, halogen, lower alkyl, -CON(R⁴)₂, guanidine, amidine,
-H, and perhaloalkyl;

E is selected from the group consisting of -H, halogen, lower alkylthio,
lower perhaloalkyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, -CN,
and -NR⁷₂;


20 X is selected from the group consisting of alkylamino, alkyl, alkenyl,
alkynyl, alkyl(carboxyl), alkyl(hydroxy), alkyl(phosphonate), alkyl(sulfonate),
aryl, alkylaminoalkyl, alkoxyalkyl, alkylthioalkyl, alkylthio, alicyclic, 1,1-
dihaloalkyl, carbonylalkyl, aminocarbonylamino, alkylaminocarbonyl,
alkylcarbonylamino, aralkyl, and alkylaryl, all optionally substituted; or together
25 with Y forms a cyclic group including cyclic alkyl, heterocyclic, and aryl;

Y is selected from the group consisting of -H, alkyl, alkenyl, alkynyl, aryl,
alicyclic, aralkyl, aryloxyalkyl, alkoxyalkyl, -C(O)R³, -S(O)₂R³, -C(O)-OR³,
-CONHR³, -NR²₂, and -OR³, all except H are optionally substituted; or together
30 with X forms a cyclic group including aryl, cyclic alkyl, and heterocyclic;

R¹ is independently selected from the group consisting of
-H, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or
thiocarbonate, -C(R²)₂-aryl, alkylaryl, -C(R²)₂OC(O)NR²₂,

-NR²-C(O)-R³, -C(R²)₂-OC(O)R³, C(R²)₂-O-C(O)OR³, -C(R²)₂OC(O)SR³, alkyl-S-C(O)R³, alkyl-S-S-alkylhydroxy, and alkyl-S-S-S-alkylhydroxy, or together R¹ and R¹ are -alkyl-S-S-alkyl to form a cyclic group, or together R¹ and R¹ are

5



wherein

V and W are independently selected from the group consisting of
 10 hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and -R⁹; or

together V and Z are connected to form a cyclic group containing 3-5
 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy,
 alkoxy, or aryloxy, or attached to a carbon atom that is three atoms
 15 from an oxygen attached to the phosphorus; or

together V and W are connected to form a cyclic group containing 3
 carbon atoms substituted with hydroxy, acyloxy, alkoxy, alkoxy,
 alkylthiocarboxy, hydroxymethyl, and aryloxy, or attached to a carbon atom
 that is three atoms from an oxygen attached to the phosphorus;

20 Z is selected from the group consisting of -CH₂OH, -CH₂OCOR³,
 -CH₂OC(O)SR³, -CH₂OCO₂R³, -SR³, -S(O)R³, -CH₂N₃, -CH₂NR²₂, -CH₂Ar,
 -CH(Ar)OH, -CH(CH=CR²R²)OH, -CH(C≡CR²)OH, and -R²;

with the provisos that:

a) V, Z, W are not all -H; and
 25 b) when Z is -R², then at least one of V and W is not -H or -R⁹;
 R² is selected from the group consisting of R³ and -H;
 R³ is selected from the group consisting of alkyl, aryl, alicyclic, and
 aralkyl;

R⁴ is independently selected from the group consisting of -H, lower alkyl,
 30 lower alicyclic, lower aralkyl, and lower aryl;

R^5 is selected from the group consisting of lower alkyl, lower aryl, lower aralkyl, and lower alicyclic;

R^6 is independently selected from the group consisting of -H, and lower alkyl;

5 R^7 is independently selected from the group consisting of -H, lower alkyl, lower alicyclic, lower aralkyl, lower aryl, and $-C(O)R^{10}$;

R^8 is independently selected from the group consisting of -H, lower alkyl, lower aralkyl, lower aryl, lower alicyclic, $-C(O)R^{10}$, or together they form a bidendate alkyl;

10 R^9 is selected from the group consisting of alkyl, aralkyl, and alicyclic;


R^{10} is selected from the group consisting of -H, lower alkyl, $-NH_2$, lower aryl, and lower perhaloalkyl;

15 R^{11} is selected from the group consisting of alkyl, aryl, -OH, $-NH_2$ and $-OR^3$; and

pharmaceutically acceptable prodrugs and salts thereof.

37. A method of inhibiting gluconeogenesis in animal in need thereof, comprising administering to said animal an effective amount of a compound of formula (1):

20



25 wherein

A is selected from the group consisting of $-NR^8_2$, $NHSO_2R^3$, $-OR^5$, $-SR^5$, halogen, lower alkyl, $-CON(R^4)_2$, guanidine, amidine, -H, and perhaloalkyl;


E is selected from the group consisting of -H, halogen, lower alkylthio, lower perhaloalkyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, -CN, and $-NR^7_2$;

30

X is selected from the group consisting of alkylamino, alkyl, alkenyl, alkynyl, alkyl(carboxyl), alkyl(hydroxy), alkyl(phosphonate), alkyl(sulfonate), aryl, alkylaminoalkyl, alkoxyalkyl, alkylthioalkyl, alkylthio, alicyclic, 1,1-dihaloalkyl, carbonylalkyl, aminocarbonylamino, alkylaminocarbonyl, alkylcarbonylamino, aralkyl, and alkylaryl, all optionally substituted; or together with Y forms a cyclic group including cyclic alkyl, heterocyclic, and aryl;

Y is selected from the group consisting of -H, alkyl, alkenyl, alkynyl, aryl, alicyclic, aralkyl, aryloxyalkyl, alkoxyalkyl, $-C(O)R^3$, $-S(O)_2R^3$, $-C(O)-OR^3$, $-CONHR^3$, $-NR^2_2$, and $-OR^3$, all except H are optionally substituted; or together with X forms a cyclic group including aryl, cyclic alkyl, and heterocyclic;

R^1 is independently selected from the group consisting of -H, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or thiocarbonate, $-C(R^2)_2$ -aryl, alkylaryl, $-C(R^2)_2OC(O)NR^2_2$, $-NR^2-C(O)-R^3$, $-C(R^2)_2-OC(O)R^3$, $C(R^2)_2-O-C(O)OR^3$, $-C(R^2)_2OC(O)SR^3$, alkyl-S- $C(O)R^3$, alkyl-S-S-alkylhydroxy, and alkyl-S-S-S-alkylhydroxy, or together R^1 and R^1 are -alkyl-S-S-alkyl to form a cyclic group, or together R^1 and R^1 are



20

wherein

V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or

25

together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxy, or aryloxy, or attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxy, alkoxy, alkythiocarboxy, hydroxymethyl, and aryloxy, attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;


- 5 Z is selected from the group consisting of $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOR}^3$, $-\text{CH}_2\text{OC(O)SR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{SR}^3$, $-\text{S(O)R}^3$, $-\text{CH}_2\text{N}_3$, $-\text{CH}_2\text{NR}^2_2$, $-\text{CH}_2\text{Ar}$, $-\text{CH(Ar)OH}$, $-\text{CH}(\text{CH}=\text{CR}^2\text{R}^2)\text{OH}$, $-\text{CH}(\text{C}\equiv\text{CR}^2)\text{OH}$, and $-\text{R}^2$;

with the provisos that:

- a) V, Z, W are not all -H; and
- 10 b) when Z is $-\text{R}^2$, then at least one of V and W is not -H or $-\text{R}^9$;
 R^2 is selected from the group consisting of R^3 and -H;
 R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl;
 R^4 is independently selected from the group consisting of -H, lower alkyl,
 15 lower alicyclic, lower aralkyl, and lower aryl;
 R^5 is selected from the group consisting of lower alkyl, lower aryl, lower aralkyl, and lower alicyclic;
 R^6 is independently selected from the group consisting of -H, and lower alkyl;
 20 R^7 is independently selected from the group consisting of -H, lower alkyl, lower alicyclic, lower aralkyl, lower aryl, and $-\text{C(O)R}^{10}$;
 R^8 is independently selected from the group consisting of -H, lower alkyl, lower aralkyl, lower aryl, lower alicyclic, $-\text{C(O)R}^{10}$, or together they form a bidentate alkyl;
 25 R^9 is selected from the group consisting of alkyl, aralkyl, and alicyclic;
 R^{10} is selected from the group consisting of -H, lower alkyl, $-\text{NH}_2$, lower aryl, and lower perhaloalkyl;
 R^{11} is selected from the group consisting of alkyl, aryl, -OH, $-\text{NH}_2$ and
 30 $-\text{OR}^3$; and
 pharmaceutically acceptable prodrugs and salts thereof.

38. A method of treating an animal for a disease derived from abnormally elevated insulin levels, comprising administering to said animal a
 35 therapeutically effective amount of a fructose-1,6-bisphosphatase inhibitor which binds to the AMP site of FBPase.

39. The method of claim 38 wherein said inhibitor is compound of formula (1):



5

wherein

A is selected from the group consisting of -NR⁸₂, NHSO₂R³, -OR⁵, -SR⁵, halogen, lower alkyl, -CON(R⁴)₂, guanidine, amidine, -H, and perhaloalkyl;


E is selected from the group consisting of -H, halogen, lower alkylthio, lower perhaloalkyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, -CN, and -NR⁷₂;

X is selected from the group consisting of alkylamino, alkyl, alkenyl, alkynyl, alkyl(carboxyl), alkyl(hydroxy), alkyl(phosphonate), alkyl(sulfonate), aryl, alkylaminoalkyl, alkoxyalkyl, alkylthioalkyl, alkylthio, alicyclic, 1,1-dihaloalkyl, carbonylalkyl, aminocarbonylamino, alkylaminocarbonyl, alkylcarbonylamino, aralkyl, and alkylaryl, all optionally substituted; or together with Y forms a cyclic group including cyclic alkyl, heterocyclic, and aryl;

Y is selected from the group consisting of -H, alkyl, alkenyl, alkynyl, aryl, alicyclic, aralkyl, aryloxyalkyl, alkoxyalkyl, -C(O)R³, -S(O)₂R³, -C(O)-OR³, -CONHR³, -NR²₂, and -OR³, all except H are optionally substituted; or together with X forms a cyclic group including aryl, cyclic alkyl, and heterocyclic;

R¹ is independently selected from the group consisting of -H, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or thiocarbonate, -C(R²)₂-aryl, alkylaryl, -C(R²)₂OC(O)NR²₂, -NR²-C(O)-R³, -C(R²)₂-OC(O)R³, C(R²)₂-O-C(O)OR³, -C(R²)₂OC(O)SR³, alkyl-S-C(O)R³, alkyl-S-S-alkylhydroxy, and alkyl-S-S-S-alkylhydroxy, or together R¹ and R¹ are -alkyl-S-S-alkyl to form a cyclic group, or together R¹ and R¹ are

30



wherein

V and W are independently selected from the group consisting of
 5 hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or

together V and Z are connected to form a cyclic group containing 3-5
 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy,
 alkoxycarboxy, or aryloxycarboxy attached to a carbon atom that is three atoms
 10 from an oxygen attached to the phosphorus; or

together V and W are connected to form a cyclic group containing 3
 carbon atoms substituted with hydroxy, acyloxy, alkoxycarboxy,
 alkylthiocarboxy, hydroxymethyl, and aryloxycarboxy attached to a carbon atom
 that is three atoms from an oxygen attached to the phosphorus;

15 Z is selected from the group consisting of $-\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OCOR}^3$,
 $-\text{CH}_2\text{OC(O)SR}^3$, $-\text{CH}_2\text{OCO}_2\text{R}^3$, $-\text{SR}^3$, $-\text{S(O)R}^3$, $-\text{CH}_2\text{N}_3$, $-\text{CH}_2\text{NR}^2_2$, $-\text{CH}_2\text{Ar}$,
 $-\text{CH(Ar)OH}$, $-\text{CH}(\text{CH}=\text{CR}^2\text{R}^2)\text{OH}$, $-\text{CH}(\text{C}\equiv\text{CR}^2)\text{OH}$, and $-\text{R}^2$;

with the provisos that:

- a) V, Z, W are not all $-\text{H}$; and
 20 b) when Z is $-\text{R}^2$, then at least one of V and W is not $-\text{H}$ or $-\text{R}^9$;
 R^2 is selected from the group consisting of R^3 and $-\text{H}$;
 R^3 is selected from the group consisting of alkyl, aryl, alicyclic, and
 aralkyl;

R^4 is independently selected from the group consisting of $-\text{H}$, lower alkyl,
 25 lower alicyclic, lower aralkyl, and lower aryl;

R^5 is selected from the group consisting of lower alkyl, lower aryl, lower
 aralkyl, and lower alicyclic;

R^6 is independently selected from the group consisting of $-\text{H}$, and lower
 alkyl;

30 R^7 is independently selected from the group consisting of $-\text{H}$, lower alkyl,
 lower alicyclic, lower aralkyl, lower aryl, and $-\text{C(O)R}^{10}$;

R^8 is independently selected from the group consisting of -H, lower alkyl, lower aralkyl, lower aryl, lower alicyclic, $-C(O)R^{10}$, or together they form a bidendate alkyl;

R^9 is selected from the group consisting of alkyl, aralkyl, and
5 alicyclic;


R^{10} is selected from the group consisting of -H, lower alkyl, $-NH_2$, lower aryl, and lower perhaloalkyl;

R^{11} is selected from the group consisting of alkyl, aryl, -OH, $-NH_2$ and -OR³; and
10 pharmaceutically acceptable prodrugs and salts thereof.

40. The method of claim 39 wherein said disease is atherosclerosis.

41. A method of treating an animal with excess glycogen storage
15 disease, comprising administering to said animal in need thereof a therapeutically effective amount of a fructose-1,6-bisphosphatase inhibitor which binds to the AMP site of FBPase.

42. The method of claim 41 wherein said inhibitor is a compound of
20 formula (1):



wherein


25 A is selected from the group consisting of $-NR^8$, $NHSO_2R^9$, $-OR^5$, $-SR^5$, halogen, lower alkyl, $-CON(R^4)_2$, guanidine, amidine, -H, and perhaloalkyl;

E is selected from the group consisting of -H, halogen, lower alkylthio, lower perhaloalkyl, lower alkyl, lower alkenyl, lower alkynyl, lower alkoxy, -CN,
30 and $-NR^7$;

- X is selected from the group consisting of alkylamino, alkyl, alkenyl, alkynyl, alkyl(carboxyl), alkyl(hydroxy), alkyl(phosphonate), alkyl(sulfonate), aryl, alkylaminoalkyl, alkoxyalkyl, alkylthioalkyl, alkylthio, alicyclic, 1,1-dihaloalkyl, carbonylalkyl, aminocarbonylamino, alkylaminocarbonyl, alkylcarbonylamino, aralkyl, and alkylaryl, all optionally substituted; or together with Y forms a cyclic group including cyclic alkyl, heterocyclic, and aryl;

- Y is selected from the group consisting of -H, alkyl, alkenyl, alkynyl, aryl, alicyclic, aralkyl, aryloxyalkyl, alkoxyalkyl, $-C(O)R^3$, $-S(O)_2R^3$, $-C(O)-OR^3$, $-CONHR^3$, $-NR^2_2$, and $-OR^3$, all except H are optionally substituted; or together with X forms a cyclic group including aryl, cyclic alkyl, and heterocyclic;

- R^1 is independently selected from the group consisting of -H, alkyl, aryl, alicyclic where the cyclic moiety contains a carbonate or thiocarbonate, $-C(R^2)_2$ -aryl, alkylaryl, $-C(R^2)_2OC(O)NR^2_2$, $-NR^2-C(O)-R^3$, $-C(R^2)_2-OC(O)R^3$, $C(R^2)_2-O-C(O)OR^3$, $-C(R^2)_2OC(O)SR^3$, alkyl-S- $C(O)R^3$, alkyl-S-S-alkylhydroxy, and alkyl-S-S-S-alkylhydroxy, or together R^1 and R^1 are -alkyl-S-S-alkyl to form a cyclic group, or together R^1 and R^1 are



20

wherein

- V and W are independently selected from the group consisting of hydrogen, aryl, substituted aryl, heteroaryl, substituted heteroaryl, 1-alkenyl, 1-alkynyl, and $-R^9$; or together V and Z are connected to form a cyclic group containing 3-5 atoms, optionally 1 heteroatom, substituted with hydroxy, acyloxy, alkoxy, alkoxy, or aryloxy, attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus; or

together V and W are connected to form a cyclic group containing 3 carbon atoms substituted with hydroxy, acyloxy, alkoxy, alkoxy, alkylthiocarboxy, hydroxymethyl, and aryloxy, attached to a carbon atom that is three atoms from an oxygen attached to the phosphorus;

- 5 Z is selected from the group consisting of -CH₂OH, -CH₂OCOR³, -CH₂OC(O)SR³, -CH₂OCO₂R³, -SR³, -S(O)R³, -CH₂N₃, -CH₂NR², -CH₂Ar, -CH(Ar)OH, -CH(CH=CR²R²)OH, -CH(C≡CR²)OH, and -R²;

with the provisos that:

- a) V, Z, W are not all -H; and
10 b) when Z is -R², then at least one of V and W is not -H or -R³;
R² is selected from the group consisting of R³ and -H;
R³ is selected from the group consisting of alkyl, aryl, alicyclic, and aralkyl;

- R⁴ is independently selected from the group consisting of -H, lower alkyl,
15 lower alicyclic, lower aralkyl, and lower aryl;

R⁵ is selected from the group consisting of lower alkyl, lower aryl, lower aralkyl, and lower alicyclic;

R⁶ is independently selected from the group consisting of -H, and lower alkyl;

- 20 R⁷ is independently selected from the group consisting of -H, lower alkyl, lower alicyclic, lower aralkyl, lower aryl, and -C(O)R¹⁰;

R⁸ is independently selected from the group consisting of -H, lower alkyl, lower aralkyl, lower aryl, lower alicyclic, -C(O)R¹⁰, or together they form a bidentate alkyl;

- 25 R⁹ is selected from the group consisting of alkyl, aralkyl, and alicyclic;

R¹⁰ is selected from the group consisting of -H, lower alkyl, -NH₂, lower aryl, and lower perhaloalkyl;

- 30 R¹¹ is selected from the group consisting of alkyl, aryl, -OH, -NH₂ and -OR³; and

pharmaceutically acceptable prodrugs and salts thereof.

43. The method of claims 34, 35, 36, 37, 38, 39, 40, 41, or 42 wherein said compound is administered orally.

1/10




FIGURE 1

2/10


In Vitro inhibition of hlFBPase

FIGURE 2

3/10

Displacement of AMP from hlFBPase




FIGURE 3

4/10

Effect of Compound 2.7 on Gluconeogenesis from Dihydroxyacetone in Rat Hepatocytes




FIGURE 4A






FIGURE 4B

Inhibition of Glucose Production From Lactate Pyruvate (Rat Hepatocytes)

**FIGURE 5**

6/10

Compound 16.4 in 18h-Fasted, Normal Rats (i.p.)**FIGURE 6**

Compound 2.7 in 18-hour fasted rats
(20 mgs/kg, i.p.)




FIGURE 7




FIGURE 8

8/10

24h fasted ZDF Rats + COMPOUND 2.7




FIGURE 9A





FIGURE 9B

9/10

**FIGURE 10**

Rat Hepatocytes: Inhibition of Glucose Production and Cellular Penetration




FIGURE 11B




FIGURE 11A

INTERNATIONAL SEARCH REPORT

International Application No

PC1/US 98/04502

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07F9/6561 A61K31/675

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07F A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 427 799 B (GENSIA PHARMACEUTICALS, INC.) 30 November 1994 cited in the application see the whole document ---	1-43
Y	EP 0 354 322 A (AMERICAN CYANAMID CO.) 14 February 1990 see the whole document ---	1-43
Y	WO 94 07867 A (PFIZER INC.) 14 April 1994 see the whole document ---	1-43
Y	EP 0 753 525 A (NIPPON SHINYAKU CO., LTD.) 15 January 1997 see pages 28 and 44, examples 5 and 6; claims --- -/-	1-43

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

3 June 1998

Date of mailing of the international search report

19. 06. 1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Beslier, L

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/04502

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 632 048 A (MITSUBISHI KASEI CORP.) 4 January 1995 cited in the application see the whole document ---	11-43
A	WO 94 22864 A (STERLING WINTHROP INC.) 13 October 1994 see the whole document -----	1-33

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 98/04502

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 33-43 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/04502

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 427799 B	22-05-1991	US 5082829 A US 5200525 A DE 69014562 D DE 69014562 T EP 0427799 A AT 114474 T AU 5086190 A AU 5236093 A CA 2008325 A DK 427799 T IE 75372 B JP 3504728 T WO 9009163 A US 5658889 A	21-01-1992 06-04-1993 12-01-1995 22-06-1995 22-05-1991 15-12-1994 05-09-1990 24-02-1994 24-07-1990 08-05-1995 10-09-1997 17-10-1991 23-08-1990 19-08-1997
EP 354322 A	14-02-1990	US 4943629 A JP 2091091 A	24-07-1990 30-03-1990
WO 9407867 A	14-04-1994	AU 683620 B AU 4669793 A CA 2145640 A EP 0662962 A FI 934224 A HU 65531 A JP 7507072 T NO 951155 A NZ 254550 A US 5728704 A ZA 9307142 A	20-11-1997 26-04-1994 14-04-1994 19-07-1995 29-03-1994 28-06-1994 03-08-1995 26-05-1995 22-08-1997 17-03-1998 23-03-1995
EP 753525 A	15-01-1997	WO 9526971 A	12-10-1995
EP 632048 A	04-01-1995	CA 2126601 A CN 1106017 A JP 7070159 A	30-12-1994 02-08-1995 14-03-1995
WO 9422864 A	13-10-1994	AT 155467 T AU 6449394 A CA 2159631 A DE 69404289 D	15-08-1997 24-10-1994 13-10-1994 21-08-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/04502

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9422864 A		DE 69404289 T	19-02-1998
		DK 691968 T	23-02-1998
		EP 0691968 A	17-01-1996
		ES 2107205 T	16-11-1997
		JP 8508492 T	10-09-1996
		US 5576427 A	19-11-1996
